

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 February 2003 (06.02.2003)

PCT

(10) International Publication Number
WO 03/010202 A1

(51) International Patent Classification⁷: **C07K 16/46**

(21) International Application Number: **PCT/KR02/01427**

(22) International Filing Date: **26 July 2002 (26.07.2002)**

(25) Filing Language: **Korean**

(26) Publication Language: **English**

(30) Priority Data:
2001-45028 26 July 2001 (26.07.2001) KR

(71) Applicant (for all designated States except US): **MEDEX-GEN CO. LTD.** [KR/KR]; 2th Floor, Medical Bldg A, Hanyang University College of Medicine, 17 Haengdang-dong, Seongdong-gu, 133-791 Seoul (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHUNG,**

Yong-Hoon [KR/KR]; #405-804 Jugong Apt., Dun-chon-dong, Gandong-gu, 134-060 Seoul (KR). **HAN, Ji-Woong** [KR/KR]; #201 HanYang Villa 24-5, Gueui-2dong, Gwangjin-gu, 143-816 Seoul (KR). **LEE, Hye-Ja** [KR/KR]; #607 ChungSil Apt., Gae-bonghon-dong, Guro-gu, 152-806 Seoul (KR). **CHOI, Eun-Yong** [KR/KR]; 19-1 Chungchun-1dong, Pupy-ong-gu, 403-854 Inchun-si (KR). **KIM, Jin-Mi** [KR/KR]; 409-287 Shillimbong-dong, Gwanak-gu, 151-029 Seoul (KR).


(74) Agents: **LEE, Sei-Jin** et al.; 17th Floor, City Air Tower, 159-9 Samsung-dong, Gangnam-gu, 135-973 Seoul (KR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,


[Continued on next page]

(54) Title: **CONCATAMERIC IMMUNOADHESION**

a



b



(57) Abstract: Disclosed are concatameric proteins comprising two soluble domains, in which the C-terminus of a soluble domain of a biologically active protein is linked to the N-terminus of an identical soluble domain or a distinct soluble domain of a biologically active protein. Also, the present invention disclosed dimeric proteins formed by formation of intermolecular disulfide bonds at the hinge region of two monomeric proteins formed by linkage of a concatamer of two identical soluble extracellular regions of proteins involving immune response to an Fc fragment of an immunoglobulin molecule, their glycosylated proteins, DNA constructs encoding the monomeric proteins, recombinant expression plasmids containing the DNA constructs, host cells transformed or transfected with the recombinant expression plasmids, and a method of preparing the dimeric proteins by culturing the host cells. Further, the present invention disclosed pharmaceutical or diagnostic compositions comprising the dimeric protein or its glycosylated form.

WO 03/010202 A1



SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

CONCATAMERIC IMMUNOADHESION

TECHNICAL FIELD

The present invention relates to concatameric proteins, and more specifically, concatamerized structure of biologically active protein domains where C-terminal end of extracellular soluble domain of biologically active protein is fused to N-terminal end of the same or other extracellular soluble domain of biologically active protein, and dimerization of two concatamers by coupling to hinge region of Fc fragment of immunoglobulin, and glycosylated forms of the concatameric proteins.

BACKGROUND ART

The activity of cytokine is associated with pathologic severity of inflammatory and /or immune response to various antigenic stimulations. Many antigen specific antibodies and soluble receptors which could recognize cytokines are currently in use to inhibit the function of cytokines for the therapeutic purposes (WO 93/016184, WO 96/02576, WO 96/023067, WO 1997/03682, and US 5,434,131, 5,656,272, 5,977,318, 6,210,661, 6,225,117). Antibodies and soluble receptors inhibit cytokine signal transduction by disturbing interaction between cytokines and their receptors on cell surface.

Soluble receptors used as functional inhibitors of cytokine that fused to heavy chains of human immunoglobulins were disclosed by Capon et al. (Nature 337:5254, 1989), and thereafter many patents were disclosed inventions related to fusion proteins of soluble receptors and immunoglobulins (US patent 5,521,288, 5,844,095, 6,046,310, 6,090,914, 6,100,383, 6,225,448).

Generally, fusion proteins of soluble receptors and immunoglobulins have following advantages (Capon et al., Nature 337:5254, 1989)

1. Increase in total avidity to ligand by forming bivalency via dimerization.
2. Increase in blood half-life of proteins, that is, increase in molecular stability
3. Activation of effector cells by Fc fragment of immunoglobulin heavy chain
4. Convenience of purification by using affinity column, e.g. using protein A

Most fusion proteins of receptor extracellular domain and immunoglobulin heavy chain are composed of heavy chain without CH1 domain, which result in dimers not binding to light chains. This structure is more desirable for the function of proteins and receptors involving immune response. For example, TNFR(WO92/16221, WO95/34326)-immunoglobulin fusion proteins disclosed in WO94/06476 and US 5,447,851 have been used for the inhibition of TNF-mediated inflammation. It is well known that TNFR-immunoglobulin fusion proteins have a higher affinity than original monomeric molecules (Lesslauer et al., Eur. J. Immunol. 21:2883, 1991; Ashkenazi et al., Proc. Natl. Acad. Sci. 88:10535, 1991; Peppe et al., J. Exp. Med. 174:1483, 1991; Mohler et al., J. Immunol. 151:1548, 1993).

For the improved inhibition of TNF mediated response, one can increase efficacy by multimerizing soluble extracellular domains of TNFR, CD2, and CTLA-4. For example, when fusion proteins of TNFR's extracellular domains bound with immunoglobulin heavy chain(heavy chain fusion protein) and with light chain(light chain fusion protein) respectively are coexpressed in the same cell, one can produce fusion proteins as a tetrameric form by linking heavy chain to heavy and light chains. This tetramer showed much more increased efficacy than monomeric or dimeric forms as presented by Scallon et al. (Cytokine 7:759, 1995).

However, this method had many difficulties for commercialization such as simultaneous expression of two different fusion genes in the same cell line, remarkably lower production yields of multimeric form, and difficulty in purifying multimeric high

molecular weight forms. For these reasons, immunoglobulin fusion proteins currently in use are only heavy chain fused form.

Therefore, there is considerable demand for the development of methods of producing multimeric protein therapeutics with high yield and efficient purification procedures.

DISCLOSURE OF INVENTION

5 The present inventors have manufactured concatameric proteins by fusing the C-terminal end of soluble domain of biologically active protein to the N-terminal end of soluble domain of the same or other biologically active protein by using DNA recombination techniques. Also, the present inventors have dimerized this concatamers by linking it to the hinge region of Fc fragment of immunoglobulin and added more
10 glycosylations by using DNA mutagenesis techniques. And the present inventors have found that concatamerized protein dimers and their glycosylated forms show increased efficacy and stability compared to conventional monomeric fusion proteins.

 Therefore, in one aspect, the present invention provides concatameric proteins where C-terminal end of soluble domain of biologically active proteins is fused to N-
15 terminal end of soluble domain of the same or other biologically active proteins.

 In another aspect, the present invention provides dimeric proteins formed by disulfide bond at hinge region of two monomeric proteins whose concatamerized part is fused to hinge region of Fc fragment of immunoglobulin.

 Also in another aspect, the present invention provides DNA constructs that
20 encode monomeric fusion proteins whose concatamerized domain is fused to hinge region of Fc fragment of immunoglobulins.

Also in another aspect, the present invention provides DNA plasmids comprising a DNA construct that encodes monomeric fusion protein whose concatamerized part is fused to hinge region of Fc fragment of immunoglobulin.

5 Also in another aspect, the present invention provides host cells transfected or transformed with recombinant DNA plasmids including a DNA construct that encodes monomeric fusion protein whose concatamerized part is fused to hinge region of Fc fragment of immunoglobulin.

10 Also in another aspect, the present invention provides a method for culturing the host cells, which were transfected or transformed with recombinant DNA plasmids including a DNA construct that encodes monomeric fusion protein whose concatamerized part is fused to hinge region of Fc fragment of immunoglobulin, under culture condition for expression of DNA constructs encoding concatameric fusion protein coupled to hinge region of Fc fragment of immunoglobulin, and manufacturing dimeric concatamers formed by disulfide bond at hinge region of two monomeric concatamers described as above
15 including the process of purification of the proteins described as above from cell culture.

Also in another aspect, the present invention provides a method for culturing the host cells, which were transfected or transformed with recombinant DNA plasmids including a DNA construct that encodes monomeric fusion protein whose concatamerized part of immunomodulatory function is fused to hinge region of Fc fragment of immunoglobulin and is inserted with glycosylation motifs, under the best condition which
20 is suitable for expression of DNA constructs that encode monomeric fusion protein whose concatamerized part of immune function is fused to hinge region of Fc fragment of immunoglobulin, and for manufacturing glycosylated dimers formed by disulfide bond at hinge region of two monomeric proteins described as above including the process of
25 purification of the glycosylated proteins described as above from cell culture.

Also in another aspect, the present invention provides DNA primers for inserting glycosylation motif into the DNA constructs that encode monomeric fusion

proteins whose concatamerized part is fused to hinge region of Fc fragment of immunoglobulins.

Also in another aspect, the present invention provides the glycosylated dimers formed by disulfide bond at hinge region of two monomeric proteins whose concatamerized part involving immune response is fused to hinge region of Fc fragment of immunoglobulins.

Also in another aspect, the present invention provides the pharmaceutical compositions comprising dimers formed by disulfide bond at hinge region of two monomeric proteins whose concatamerized part involving immune response is fused to hinge region of Fc fragment of immunoglobulins in a pharmaceutically effective amount and in a pharmaceutically acceptable carrier.

Also in another aspect, the present invention provides the pharmaceutical compositions comprising glycosylated dimers formed by disulfide bond at hinge region of two monomeric proteins whose concatamerized part involving immune response is fused to hinge region of Fc fragment of immunoglobulins in a pharmaceutically effective amount and in a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and other objects, features and other advantages of the present invention will be more clearly understood from the following detailed description taken in conjunction with the accompanying drawings, in which:

Fig. 1 is a schematic view showing a process of preparing a DNA construct encoding a conventional simple fusion monomeric protein through polymerase chain reaction (PCR);

Fig. 2 is a schematic view showing a process of preparing a DNA construct encoding a concatameric fusion monomeric protein according to the present invention through PCR;

Fig. 3a shows structures of [TNFR/Fc]₂, [CD2/Fc]₂ or [CTLA4/Fc]₂ fusion proteins, which are simple fusion dimeric proteins formed through homodimerization in cells of TNFR/Fc, CD2/Fc or CTLA4/Fc fusion proteins as examples of conventional simple fusion monomeric proteins;

5 Fig. 3b shows structures of [TNFR-TNFR/Fc]₂, [CD2-CD2/Fc]₂ or [CTLA4-CTLA4/Fc]₂ fusion proteins, which are concatameric fusion dimeric proteins formed through homodimerization in cells of TNFR-TNFR/Fc, CD2-CD2/Fc or CTLA4-CTLA4/Fc fusion proteins as embodiments of the concatameric fusion dimeric protein according to the present invention;

10 Fig. 4a shows a structure of [TNFR1-TNFR1/Fc]₂, as an embodiment of a concatameric fusion dimeric protein according to the present invention;

Fig. 4b shows a structure of [TNFR2-TNFR2/Fc]₂, as another embodiment of the concatameric fusion dimeric protein according to the present invention;

15 Fig. 4c shows a structure of [CD2-CD2/Fc]₂, as a further embodiment of the concatameric fusion dimeric protein according to the present invention;

Fig. 4d shows a structure of [CTLA4-CTLA4/Fc]₂, as a still further embodiment of the concatameric fusion dimeric protein according to the present invention;

20 Fig. 5 is a diagram showing a process of constructing a recombinant expression plasmid pTR11Ig-Top10' expressing a concatameric fusion monomeric protein TNFR1-TNFR1/Fc according to the present invention;

Fig. 6 is a diagram showing a process of constructing a recombinant expression plasmid pCD22Ig expressing a concatameric fusion monomeric protein CD2-CD2/Fc according to the present invention;

25 Fig. 7 is a map of a recombinant expression plasmid pTR11Ig-Top10' expressing a concatameric fusion monomeric protein TNFR1-TNFR1/Fc according to the present invention;

Fig. 8 is a map of a recombinant expression plasmid pTR22Ig-Top10' expressing a concatameric fusion monomeric protein TNFR1-TNFR1/Fc according to the present invention;

30 Fig. 9 is a map of a recombinant expression plasmid pCD22Ig expressing a concatameric fusion monomeric protein CD2-CD2/Fc according to the present invention;

Fig. 10 is a map of a recombinant expression plasmid pCT44Ig expressing a concatameric fusion monomeric protein CTLA4-CTLA4/Fc according to the present invention;

5 Fig. 11 is a map of a recombinant expression plasmid pTR11Ig-MG expressing a concatameric fusion monomeric protein mgTNFR1-TNFR1/Fc containing four glycosylation motif peptides according to the present invention;

Fig. 12 is a map of a recombinant expression plasmid pTR22Ig-MG expressing a concatameric fusion monomeric protein mgTNFR2-TNFR2/Fc containing two glycosylation motif peptides according to the present invention;

10 Fig. 13 is a map of a recombinant expression plasmid pCD22Ig-MG expressing a concatameric fusion monomeric protein mgCD2-CD2/Fc containing two glycosylation motif peptides according to the present invention;

Fig. 14 is a map of a recombinant expression plasmid pCT44Ig-MG expressing a concatameric fusion monomeric protein mgCTLA4-CTLA4/Fc containing three glycosylation motif peptides according to the present invention;

15 Fig. 15 shows a result of SDS-PAGE of purified concatameric fusion dimeric proteins [TNFR1-TNFR1/Fc]₂ and [TNFR2-TNFR2/Fc]₂ under reducing or non-reducing conditions;

Fig. 16 is a graph showing inhibitory effect of the conventional simple fusion dimeric proteins [TNFR1/Fc]₂(●) and [TNFR2/Fc]₂(○) and the concatameric fusion dimeric proteins [TNFR1-RNFR1/Fc]₂(▼) and [TNFR2-TR2Fc]₂(▽) according to the present invention against cytotoxic activity of TNF-alpha;

20 Fig. 17 is a graph showing inhibitory effect of the conventional simple fusion dimeric proteins [TNFR1/Fc]₂(●) and [TNFR2/Fc]₂(○) and the concatameric fusion dimeric proteins [TNFR1-RNFR1/Fc]₂(▼) and [TNFR2-TR2Fc]₂(▽) according to the present invention against cytotoxic activity of TNF-beta;

25 Fig. 18 is a graph showing inhibitory effect of the conventional simple fusion dimeric protein [CD2/Fc]₂(●), the known immunosuppressive agent cyclosporin A (▼) and the concatameric fusion dimeric protein [CD2-CD2/Fc]₂(○) according to the present invention on the proliferation of active T lymphocytes;

30 Fig. 19 is a graph showing inhibitory effect of the conventional simple fusion

dimeric protein [CTLA4/Fc]₂(●), the known immunosuppressive agent cyclosporin A (▼) and the concatameric fusion dimeric protein [CTLA4-CTLA4/Fc]₂ (○) according to the present invention on the proliferation of active T lymphocytes;

5 Fig. 20 is a graph showing blood half-life of the conventional simple fusion dimeric protein [TNFR1/Fc]₂(●), the concatameric dimeric protein [TNFR1-TNFR1/Fc]₂ (○) and a glycosylated concatameric fusion dimeric protein [mgTNFR1-TNFR1/Fc]₂ (▽) according to the present invention;

10 Fig. 21 is a graph showing blood half-life of the conventional simple fusion dimeric protein [CD2/Fc]₂(●), the concatameric fusion dimeric protein [CD2-CD2/Fc]₂ (○) and a glycosylated concatameric fusion dimeric protein [mgCD2-CD2/Fc]₂ (▽) according to the present invention;

15 Fig. 22 is a graph showing blood half-life of the conventional simple fusion dimeric protein [CTLA4/Fc]₂(●), the concatameric fusion dimeric protein [CTLA4-CTLA4/Fc]₂ (○) and a glycosylated concatameric fusion dimeric protein [mgCTLA4-CTLA4/Fc]₂ (▽) according to the present invention; and

20 Fig. 23 is a graph showing inhibitory effect of PBS (●) as a control, the conventional simple fusion dimeric proteins [TNFR1/Fc]₂(■) and [TNFR2/Fc]₂(▲), and concatameric fusion dimeric proteins [TNFR1-TNFR1/Fc]₂ (×) and [TNFR2-TNFR2/Fc]₂ (△) according to the present invention on the induction of collagen-induced arthritis (CIA) in DBA/1 mice.

BEST MODE FOR CARRYING OUT THE INVENTION

25 The present invention is generally directed to concatameric proteins, and more particularly, to immunoadhesion molecules. Immunoadhesion molecules are typically formed by fusion of the Fc fragment of immunoglobulin (Ig) to a ligand-binding region of a receptor or an adhesion molecule, and thus have a structure similar to that of an antibody. The typical immunoadhesion molecules known in the art have a structure of an antibody in which the variable region is substituted with a ligand-binding region of a receptor while retaining the Fc fragment. A wide variety of immunoadhesion molecules
30 are suggested in the literature. However, immunoadhesion molecules according to the

present invention have different structure with the conventional immunoadhesion molecules, and there is also no prior art predicting or describing preparation of the immunoadhesion molecules according to the present invention.

Definition of Terms

5 For full understanding of the characteristic structure of the immunoadhesion molecules according to the present invention, exact definitions of the terms used in the present invention are given as follows. In general, all of the technical and scientific terms being not additionally defined in the present invention have meanings commonly used in the art. However, although having meanings commonly used in the art, the
10 following terms are defined to give a clearer understanding of their meanings and make the scope of the present invention clear, as follows.

The term "immunoglobulin", as used herein, refers to protein molecules being produced in B cells and serving as antigen receptors specifically recognizing a wide variety of antigens. The molecules have a Y-shaped structure consisting of two identical
15 light chains (L chains) and two identical heavy chains (H chains), in which the four chains are held together by a number of disulfide bonds, including the disulfide bridge between the H chains at the hinge region. The L and H chains comprise variable and constant regions. The L chain variable region associates with the H chain variable region, thus producing two identical antigen-binding regions. According to features of the constant
20 regions of H chains, immunoglobulins (Ig) are classified into five isotypes, A (IgA), D (IgD), E (IgE), G (IgG) and M (IgM). Each subtype possesses unique structural and biological properties. For example, IgG has slightly different Fc structure, compared with other isotypes. In addition, IgG and IgA have a number of subtypes. For example, the human IgG isotype has four subtypes, IgG1, IgG2, IgG3 and IgG4, which
25 have $\gamma 1$, $\gamma 2$, $\gamma 3$ and $\gamma 4$ H chains, respectively. Biological functions of immunoglobulin molecules, such as complement activation, Fc receptor-mediated phagocytosis and antigen-dependent cytotoxicity, are mediated by structural determinants (complementarity-determining regions) in the Fc region of H chains. Such an Fc region of H chains is used for construction of dimeric proteins according to the present

invention, and may be derived from all isotypes and subtypes of immunoglobulin as described above.

5 The term "Fc fragment of an immunoglobulin molecule", as used herein, refers to a fragment having no antigen-binding activity and being easily crystallized, which comprises a hinge region and CH2 and CH3 domains, and a portion responsible for binding of an antibody to effector materials and cells. Therefore, the Fc fragment mentioned in the present invention can be different from that described in some literatures, but includes the hinge region. Such description of the Fc fragment is given to supply convenience in describing the present invention, and will be fully understood by those of ordinary skill in the art with reference to the specification of the present invention and the accompanying drawings.

10 The term "biologically active protein", as used herein, refers to a protein, peptide or polypeptide having generally physiological or pharmaceutical activities, which retains a part of its native activities after forming a concatamer or immunoadhesion molecule. The term "biological activity", as used herein, is not limited in meaning to physiological or pharmaceutical activities. For example, some concatamers, such as those containing an enzyme can catalyze a reaction in an organic solvent. Similarly, some high-molecular weight fusion molecules containing concanavalin A or an immunoglobulin molecule are useful as diagnostic agents in laboratories.

20 Non-limiting examples of the protein, peptide or polypeptide include hemoglobin, serum proteins (e.g., blood factors including factor VII, VIII and factor IX), immunoglobulin, cytokines (e.g., interleukin), α -, β - and γ -interferon, colony-stimulating agent (e.g., G-CSF and GM-CSF), platelet-derived growth factor (PDGF), and phospholipase activating proteins (PLAPs). Other typical biological or therapeutic proteins include insulin, plant proteins (e.g., lectin and ricin), tumor necrosis factor (TNF) and its related alleles, growth factors (e.g., tissue growth factors and endothelial growth factors such as TGF α or TGF β), hormones (e.g., follicle-stimulating hormone, thyroid-stimulating hormone, antidiuretic hormone, pigment-concentrating or dispersing hormones and parathyroid hormone, luteinizing hormone-releasing hormone and its derivatives, calcitonin, calcitonin gene related peptide (CGRP), synthetic enkephalin, somatomedin, erythropoietin, hypothalamus releasing factors, prolactin, chronic gonadotrophin, tissue

plasminogen-activating agents, growth hormone-releasing peptide (GHRP), and thymic humoral factor (THF). The immunoglobulins include IgG, IgE, IgM, IgA, IgD and fragments thereof. Some proteins such as interleukin, interferon or colony-stimulating factor may be produced in a non-glycosylated form using DNA recombinant techniques.

5 The non-glycosylated proteins may be useful as biologically active materials in the present invention.

In addition, the biologically active materials useful in the present invention include any polypeptide, which has bioactivity in vivo. Examples of the biologically active materials include peptides or polypeptides, fragments of an antibody, single chain-binding proteins (see U.S. Pat. No. 4,946,778), binding molecules including fusion polypeptides of antibodies or their fragments, polyclonal antibodies, monoclonal antibodies, and catalytic antibodies. Other examples of the biologically active materials include allergen proteins, such as ragweed, antigen E, honeybee venom, or allergen of mites.

15 In addition, the biologically active material useful in the present invention includes enzymes. Examples of the enzymes include carbohydrate-specific enzymes, proteinases, oxidoreductases, transferases, hydrolases, lyases, isomerases, and ligases. In detail, non-limiting examples of the enzymes include asparaginase, arginase, arginine deaminase, adenosine deaminase, peroxide dismutase, endotoxinase, catalase, chymotrypsin, lipase, uricase, adenosine dephosphatase, tyrosinase, and bilirubin oxidase.

20 Examples of the carbohydrate-specific enzymes include glucose oxidase, glucodase, galactosidase, glucocerebrosidase, and glucouronidase.

The term "proteins involving immune response", as used herein, refers to all proteins mediating cell-to-cell signal transduction during cellular or humoral immune response and thus activating or suppressing immune response. Immunity is a process of protecting "self" from "non-self" such as bacteria or viruses. Immune response is largely divided into cellular and humoral immune response, where T and B lymphocytes play the most important role. T cells, mainly mediating cellular immune response, directly attack and kill virus-infected cells or tumor cells, or help other immune cells by secreting cytokines functioning to induce or activate immune response or inflammation.

30 B cells produce antibodies against non-self foreign materials (antigens) that enter a body,

such as bacteria or viruses, and such immune response is called cellular immune response. Cell-to-cell signal transduction is an essential process in both cellular and humoral immune responses, in which a signal molecule, that is, a ligand, interacts with a cell surface receptor acting to transduce a specific signal into a cell.

5 Representative examples of the proteins involving the immune response according to the present invention include cytokines, cytokine receptors, adhesion molecules, tumor necrosis factor receptor (TNFR), enzymes, receptor tyrosine kinases, chemokine receptors, other cell surface proteins, and soluble ligands. Non-limiting
10 examples of the cytokines include IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12, IL-17, TNF, TGF, IFN, GM-CSF, G-CSF, EPO, TPO, and M-CSF. Examples of the cytokine receptors, but are not limited to, include growth hormone receptors (GHRs), IL-13R, IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-9R, IL-15R, TNFR, TGFR, IFNR (e.g., IFN- γ R α -chain and IFN- γ R β -chain), interferon- α R, - β R and - γ R, GM-CSFR, G-CSFR, EPOR, cMpl, gp130, and Fas (Apo 1). Non-limiting examples of the
15 enzymes include influenza C hemagglutinin esterase and urokinase. The chemokine receptors are exemplified by CCR1 and CXCR1-4. Examples of the receptor tyrosine kinases, but are not limited to, include TrkA, TrkB, TrkC, Htk, REK7, Rse/Tyro-3, hepatocyte growth factor R, platelet-derived growth factor R, and Flt-1. Examples of other cell surface proteins includes CD2, CD4, CD5, CD6, CD22, CD27, CD28, CD30,
20 CD31, CD40, CD44, CD100, CD137, CD150, LAG-3, B7, B61, β -neurexin, CTLA-4, ICOS, ICAM-1, complement R-2 (CD21), IgER, lysosomal membrane gp-1, α 2-microglobulin receptor-related proteins, and sodium-releasing peptide R. Non-limiting examples of the soluble ligands include IL-10, heregulin, and keratinocyte growth factors.

25 Ligands for the proteins involving immune response according to the present invention and use thereof are well known to those of ordinary skill in the art, as summarized in Tables 1 to 7, below.

TABLE 1
Proteins involving immune response: Adhesion molecules

Adhesion molecules	Ligands	Uses
CD4	HIV gp120	Inhibition of in vivo HIV infection; and identification of CD4 domain participating in ligand binding
L-Selectin	GlyCAM-1, CD34	Prevention of neutrophil-mediated lung damage; determination of position in tissues of a ligand by histochemical staining; and isolation and cloning of ligands and determination of their properties
E-Selectin	Sialyl Lewis ^x	Prevention of neutrophil-mediated lung damage; and determination of thermodynamic properties in ligand-binding
P-Selectin	Sialyl Lewis ^x	Prevention of neutrophil-mediated lung damage; and study of functions of individual of amino acid residues in binding to cell surface
ICAM-1	CD11a/CD18	Phagocytosis of erythrocytes in malaria; inhibition of infection with rhinovirus; and anti-inflammation in diabetes
ICAM-2	CD11a/CD18	Study of activation of T cells mediated by T cell receptor
ICAM-3	CD11a/CD18	Identification of receptor domains binding to a ligand
VCAM-1	VLA-4	Study of role of VLA-4 in T lymphocyte migration to dermal inflammation sites
LFA-3	CD2	Study of role of CD2 in costimulation of T cells
L1 glycoprotein	Fibroblast growth factor receptor	Stimulation of nerve reproduction after repair; and functional comparison with FGF

TABLE 2
Proteins involving immune response: Enzymes

Enzymes	Ligands	Uses
Influenza C hemagglutinin esterase	9-0-acetylated sialic acid	Inactive enzyme used in study of tissue-specific expression of ligands
Urokinase	Urokinase receptor	Inactive enzyme developed to inhibit cancer metastasis by disturbing urokinase activation

5

TABLE 3
Proteins involving immune response: Cytokine receptors

Cytokine receptors	Ligands	Uses
IFN- γ R α -chain	IFN- γ	Inhibition of IFN-mediated autoimmunity
IFN- γ R β -chain	IFN- γ	Study of structure of subunits of a ligand-receptor complex
IL1R	IL-1	Inhibition of IL-1-mediated inflammation
IL4R	IL-4	Identification of receptor domains participating in ligand binding
Erythropoietin R	Erythropoietin	Map design of epitopes of anti-ligand antibodies
cMpl	Thrombopoietin	Isolation and cloning of ligands
gp130	IL-6-IL6R complex	Study of structure of subunits of a ligand-receptor complex

TABLE 4

Proteins involving immune response: Tumor necrosis factor receptors

TNF receptors	Ligands	Uses
TNF R-1	TNF, lymphotoxin- α	Treatment of septic shock, rheumatoid arthritis and other inflammatory diseases; and identification of domains participating in ligand binding
TNF R-2	TNF, lymphotoxin- α	Inhibition of TNF-enriched HIV replication; and prevention of collagen-induced arthritis in mice
Lymphotoxin- β R	Lymphotoxin- β	Study of structure of subunits of cell surface lymphotoxin- β
Fas/Apo-1/CD95	Fas/Apo-1/CD95 ligand	Treatment of excessive apoptosis and related diseases (e.g., AIDS); and resistance to apoptosis of lymphocytes and peripheral immune tolerance; roles of Fas ligand in T cell-mediated cytotoxicity; and isolation and cloning of ligands
CD27	CD27 ligand	Isolation and cloning of ligands
CD30	CD30 ligand	Isolation and cloning of ligands
CD40	gp39	Isolation and cloning of ligands
4-1BB	4-1BB ligand	Identification of tissues containing ligands by histochemical staining; isolation and cloning of ligands; and Study of structural determinant of potential ligand
OX40	gp34	Isolation and cloning of ligands

TABLE 5

Proteins involving immune response: Receptor tyrosine kinases

Receptor tyrosine kinases	Ligands	Uses
TrkA, B, C	Neutropin	Determination of properties of neutropin binding
Htk	Htk ligand	Isolation and cloning of ligands
REK7	AL-1	Isolation and cloning of ligands
Rse/Tyro-3	Protein S, Gas6	Identification of ligands and determination of their properties
Hepatocyte growth factor R	Hepatocyte growth factor	Identification of receptor domains participating in ligand binding
Platelet-derived growth factor R	Platelet-derived growth factor	Identification of receptor domains participating in ligand binding
Flt-1	Vesicular endothelial growth factor (VEGF)	Determination of properties of ligand binding of receptors
Flk-1/KDR	VEGF	Evaluation of selectivity of receptors for VEGF versus placenta growth factor

TABLE 6
Proteins involving immune response: Other cell surface proteins

Other cell surface proteins	Ligands	Uses
B7	CD28	Study of T cell stimulation by B cells
B61	Eck	Roles of Eck in inflammation
β -neurexin	β -neurexin ligand	Determination of properties of a signal sequence from β -neurexin
CD2	LFA-3, CD48	Identification of ligands
CD5	CD5 ligand	Study of T cell stimulation by B cells
CD6	ALCAM	Study of binding activities of cloned ligands
CD22	CD45, other sialoglycoproteins	Identification of ligands; study on roles of CD22 in T-B-cell interaction; and determination of properties of binding determinants of sialo-oligo sugar ligands
CD28	B7, B7-2	Study of T cell stimulation by B cells
CD31	CD31	Identification of CD31 domains related to homotype binding
CD44	Hyaluronate	Screening of tissues containing ligands by histochemical staining; and determination of properties of structural determinants of ligands
Complement R-2 (CD21)	C3 fragment	Inhibition of reactivity of antibody to immunosuppressive and cancer therapeutic agents
CTLA-4	B7	Identification of CTLA-4 as a secondary receptor of B7
IgE R	IgE	Inhibition of mast cell-binding of IgE as therapy of allergic diseases
Lisosome membrane gp-1	LAMP-1 ligand	Design of epitope maps of anti-ligand antibodies
α 2-microglobulin receptor-bound proteins	gp330	Determination of position of ligands in tissues by histochemical staining
Sodium-releasing peptide R	Sodium-releasing peptide	Design of epitope maps of anti-ligand antibodies; and preparation of recombinant receptors for structural study

TABLE 7
Proteins involving immune response: Soluble ligands

Soluble ligands	Ligands	Uses
IL-2	IL-2R	Extension of half-life of IL-2 in the circulation system
IL-10	IL-10R	Therapy of septic shock and transplantation rejection; and extension of half-life of IL-10 in the circulation system
Heregulin	Her4/p180 ^{erbB4}	Study of signal transduction by Her4
Keratinocyte growth factor	Keratinocyte growth factor R	Determination of position of receptors by histochemical staining

- 5 The term “soluble extracellular domain”, as used herein, refers to a portion exposed to the extracellular region of an integral membrane protein penetrating the cell membrane comprising phospholipid, wherein the integral membrane protein contains one or more transmembrane domain made up predominantly of hydrophobic amino acids.

Such an extracellular domain mainly comprises hydrophilic amino acids, which are typically positioned at the surface of a folded structure of a protein, and thus is soluble in an aqueous environment. For most cell surface receptor proteins, extracellular domains serve to bind specific ligands, while intracellular domains play an important role in signal transduction.

The term "concatamer-linked", as used herein, refers to a state in which two soluble domains of biologically active proteins are linked and thus form a long polypeptide.

The term "concatameric protein", as used herein, means a concatamer-linked protein. For example, the N-terminus of a soluble extracellular domain of a protein involving immune response is linked to the C-terminus of an identical soluble extracellular domain of the protein involving immune response, wherein the C-terminus of the former soluble extracellular domain is linked to the hinge region of an Fc fragment of an immunoglobulin molecule. Thus, two identical soluble extracellular domains of a protein involving immune response form a long polypeptide.

The term "simple fusion monomeric protein", as used herein, refers to a fusion protein having a monomeric structure consisting of a single polypeptide formed by linkage of a soluble extracellular domain of a protein involving immune response to the hinge region of an Fc fragment of an immunoglobulin molecule. A simple fusion monomeric protein may be designated "protein name/Fc" for convenience in the present invention. For example, a simple fusion monomeric protein produced by linkage of an soluble extracellular domain of TNFR1 protein involving immune response to an Fc fragment of an immunoglobulin molecule is designated TNFR1/Fc. If desired, the origin of the Fc fragment may be also specified in the designation. For example, in the case that the Fc fragment is derived from IgG1, the monomeric protein is called TNFR1/IgG1Fc.

The term "simple fusion dimeric protein", as used herein, refers to a fusion protein having a dimeric structure, in which two simple fusion monomeric proteins are joined by formation of intermolecular disulfide bonds at the hinge region. Such a simple fusion dimeric protein may be designated "[protein name/Fc]₂" for convenience in the present invention. For example, when fused by formation of intermolecular disulfide

bonds at the hinge region of two simple fusion monomeric proteins produced by linkage of an soluble extracellular domain of TNFR1 protein and an Fc fragment of an immunoglobulin molecule, the resulting fusion protein having dimeric structure is designated [TNFR1/Fc]₂. In addition, the origin of the Fc fragment may be specified in the designation, if desired. For example, in the case that the Fc fragment is derived from IgG1, the dimeric protein is designated [TNFR1/IgG1Fc]₂.

The term "concatameric fusion monomeric protein", as used herein, refers to a fusion protein having a monomeric structure consisting of a single polypeptide, in which the N-terminus of a soluble extracellular domain of a protein involving immune response is linked to the C-terminus of an identical soluble extracellular domain of the protein involving immune response, wherein the C-terminus of the former soluble extracellular domain is linked to the hinge region of an Fc fragment of an immunoglobulin molecule. A concatameric fusion monomeric protein may be designated "protein name-protein name/Fc" for convenience in the present invention. For example, when an soluble extracellular domain of TNFR1 of a simple fusion monomeric protein, produced by linkage of the soluble extracellular domain of TNFR1 protein involving immune response and an Fc fragment of an immunoglobulin molecule, is linked to an identical soluble extracellular domain of TNFR1, the resulting concatameric fusion monomeric protein is designated TNFR1-TNFR1/Fc. If desired, the origin of the Fc fragment may be specified in the designation. For example, in the case that the Fc fragment is derived from IgG1, the monomeric protein is designated TNFR1-TNFR1/IgG1Fc.

The term "concatameric fusion dimeric protein", as used herein, refers to a fusion protein having a dimeric structure, in which two concatameric fusion monomeric proteins are fused by formation of intermolecular disulfide bonds at the hinge region. A concatameric fusion dimeric protein may be designated "[protein name-protein name/Fc]₂" for convenience in the present invention. For example, when two concatameric fusion monomeric proteins, each of which is produced by linkage of a TNFR1 soluble extracellular domain of a simple fusion monomeric protein to an identical soluble extracellular domain of TNFR1 protein involving immune response, are fused by formation of intermolecular disulfide bonds at the hinge region, the resulting fusion protein having dimeric structure is designated [TNFR1-TNFR1/Fc]₂, wherein the simple

fusion monomeric protein is formed by linkage of the TNFR1 soluble extracellular domain to an Fc fragment from an immunoglobulin molecule. If desired, the origin of the Fc fragment may be specified in the designation. For example, in the case that the Fc fragment is derived from IgG1, the fusion protein is designated [TNFR1-
5 TNFR1/IgG1Fc]₂.

The term "vector", as used herein, means a DNA molecule serving as a vehicle capable of stably carrying exogenous genes into host cells. For useful application, a vector should be able to replicate, have a system for introducing itself into a host cell, and possess selectable markers. The exogenous genes, for example, include, a DNA
10 construct encoding a concatameric fusion monomeric protein.

The term "recombinant expression plasmid", as used herein, refers to a circular DNA molecule carrying exogenous genes operably linked thereto to be expressed in a host cell. When introduced into a host cell, the recombinant expression plasmid has the ability to replicate regardless of host chromosomal DNA, copy itself at a high copy
15 number, and to produce heterogeneous DNA. As generally known in the art, in order to increase the expression level of a transfected gene in a host cell, the gene should be operably linked to transcription and translation regulatory sequences functional in a host cell selected as an expression system. Preferably, the expression regulation sequences and the exogenous genes may be carried in a single expression vector containing
20 bacteria-selectable markers and a replication origin. In case that eukaryotic cells are used as an expression system, the expression vector should further comprise expression markers useful in the eukaryotic host cells.

The term "operably linked", as used herein, means an arrangement of elements of a vector, in which each element is capable of performing its innate function. Therefore,
25 a control sequence operably linked to a coding sequence can influence expression of the coding sequence. A control sequence acting to induce expression of a coding sequence does not have to be adjacent to the coding sequence. For example, when an intervening sequence is present between a promoter sequence and a coding sequence, the promoter sequence may still be "operably linked" to the coding sequence.

30 Host cells used in the present invention may be prokaryotic or eukaryotic. In addition, host cells having high introduction efficiency of foreign DNA and having high

expression levels of an introduced gene may be typically used. Examples of the host cells useful in the present invention include prokaryotic and eukaryotic cells such as *E. coli*, *Pseudomonas* sp., *Bacillus* sp., *Streptomyces* sp., fungi or yeast, insect cells such as *Spodoptera frugiperda* (Sf9), animal cells such as Chinese hamster ovary cells (CHO) or mouse cells, African green monkey cells such as COS 1, COS 7, human embryonic kidney cells, BSC 1, BSC 40 or BMT 10, and tissue-cultured human cells. When cloning a DNA construct encoding the fusion protein according to the present invention, host cells are preferably animal cells. When using COS cells, since SV40 large T antigen is expressed in COS cells, a plasmid carrying a SV 40 replication origin may be present as a multicopy episome and thus allows high expression of an exogenous gene. A DNA sequence introduced into a host cell may be homogeneous or heterogeneous to the host cell, or a hybrid DNA sequence containing a homogenous or heterogeneous DNA sequence.

In order to express a DNA sequence encoding the concatameric fusion protein according to the present invention, a wide variety of combinations of host cells as an expression system and vectors may be used. Expression vectors useful for transforming eukaryotic host cells contain expression regulation sequences from, for example, SV40, bovine papillomavirus, adenovirus, adeno-associated viruses, cytomegalovirus and retroviruses. Expression vectors useful in bacterial host cells include bacterial plasmids from *E. coli*, which are exemplified by pBluescript, pGEX2T, pUC, pCR1, pBR322, pMB9 and derivatives thereof, plasmids having a broad range of host cells, such as RP4, phage DNAs, exemplified by a wide variety of λ phage derivatives including λ gt10, λ gt11 and NM989, and other DNA phages, exemplified by filamentous single-stranded DNA phages such as M13. Expression vectors useful in yeast cells include 2 μ plasmid and derivatives thereof. Expression vectors useful in insect cells include pVL 941.

The term "transformation", as used herein, means introducing DNA into a suitable host cell so that the DNA is replicable, either as an extrachromosomal element, or by chromosomal integration.

The term "transfection", as used herein, refers to the taking up of an expression vector by a suitable host cell, whether any coding sequences are in fact expressed or not.

The term "signal sequence", as used herein, means an amino acid sequence mediating transport of an expressed protein to the outside of the cell membrane, and is also

called a "leader sequence". Cell surface proteins or secretory proteins, which are transported to the outside of the cell membrane, have an N-terminal sequence typically cut by signal peptidase in the cell membrane. Such a N-terminal sequence is called a signal sequence or signal peptide, or a leader sequence or leader peptide. Secretory (or
5 transported) proteins or all proteins present outside of the cell membrane or in the extracellular environment have a specific signal sequence. There is no specific homology between such signal sequences and same proteins have different signal sequences according to their origin. Secondary structure or distribution of nonpolar and charged residues is more important for proper function of the signal sequences than primary structures thereof.
10 Although not having specific homology, the signal sequences share several common features, as follows. The signal sequences contain an N domain at their N-termini, which is a hydrophilic region comprising one or more positively charged residues, and an H-domain follows the N domain, which is a somewhat long hydrophobic region. In the case of *E. coli*, the signal sequence comprises about 18-30 amino acids. The N domain
15 contains many cationic amino acids such as Lys or Arg, and thus has a net positive charge. Many hydrophobic amino acids such as Ala or Leu are found in the H domain, and polar or charged amino acids such as Pro, Lys, Arg, Asn or Glu are rarely in the H domain. A large number of amino acids such as Ala and Leu residues form an α -helical structure to facilitate membrane penetration. A C domain is positioned between the H domain and an
20 actually secreted portion of a protein. The C domain is less hydrophobic, and contains a sequence capable of being recognized by signal peptidase such as LebB or LspA. There have been no reports about an exact site cleaved by the signal peptidase, but the signal peptidase is typically known to mostly cleave behind the Ala-X-Ala sequence in the C domain. Preproteins containing the above-mentioned signal sequence arrive at the cell
25 membrane through interaction with several proteins, and fold to their mature forms through cleavage of a specific region of a signal peptide. Such a signal sequence is very important in strategies to express a desired protein on the cell surface or in the extracellular environment. Foreign proteins and fusion proteins should be stably transported to the extracellular environment at high efficiency. Typically, cell surface proteins having
30 excellent secretory ability are useful for cell surface expression of foreign proteins or fusion

proteins, which typically have secretory signal sequences capable of offering excellent secretion efficiency.

Preparation of the concatameric fusion dimeric protein according to the present invention

5 The concatameric fusion dimeric protein according to the present invention is generally prepared by (a) preparing a DNA construct encoding a simple fusion monomeric protein using a gene encoding an Fc fragment of an immunoglobulin molecule and a gene encoding a soluble extracellular domain of a protein involving immune response; (b) inserting by polymerase chain reaction (PCR) a recognition sequence of a restriction
10 enzyme into the prepared simple fusion monomeric protein-encoding DNA construct and an identical gene to the gene encoding a soluble extracellular domain of a protein involving immune response, respectively; (c) cleaving the recognition sequence of a restriction enzyme in the simple fusion monomeric protein-coding DNA construct and the gene
15 encoding a soluble extracellular domain of a protein involving immune response using the restriction enzyme recognizing the recognition sequence; (d) ligating the cleaved DNA fragments using ligase to produce a DNA construct encoding a concatameric fusion monomeric protein (see, Fig. 2); (e) operably linking the prepared DNA construct encoding a concatameric fusion monomeric protein to a vector to produce a recombinant expression
20 plasmid; (f) transforming or transfecting a host cell with the recombinant expression plasmid; and (g) culturing the transformant or transfectant under conditions suitable for expression of the DNA construct encoding a concatameric fusion monomeric protein and then isolating and purifying a concatameric fusion dimeric protein of interest.

 A DNA fragment encoding a soluble extracellular domain of a protein involving immune response is produced by PCR using a primer containing a recognition sequence of
25 a specific restriction enzyme and a sequence encoding a leader sequence, and a primer containing an antisense sequence encoding the 3' end of the soluble extracellular domain and a portion of the 5' end of a specific region of Fc fragment of an immunoglobulin molecule.

A DNA fragment encoding a specific region of the Fc fragment of an immunoglobulin molecule is produced by PCR using a primer having a sequence encoding a portion of the 3' end of the soluble extracellular domain of the protein involving immune response and a sequence encoding the 5' end of the specific region of the Fc fragment of an immunoglobulin molecule, and another primer having an antisense sequence encoding a recognition sequence of a specific restriction enzyme and the 3' end of a specific region of the Fc fragment of an immunoglobulin molecule.

The DNA fragment encoding a soluble extracellular domain of a protein involving the immune response and the DNA fragment encoding a specific region of Fc fragment of an immunoglobulin molecule, as described above, are mixed in a test tube. After denaturation, the DNA is re-annealed. Then, a complete double-stranded DNA fragment is produced by polymerization using DNA polymerase at the 3' end of each DNA hybrid. Using the resulting double-stranded DNA fragment, another polymerase chain reaction (PCR) is carried out with the primer having a sequence encoding a soluble extracellular domain of a protein involving immune response and the primer encoding the 3' end of a specific region of the Fc fragment of an immunoglobulin molecule, in order to amplify a immunoglobulin fusion gene comprising a sequence corresponding to the DNA fragment encoding a soluble extracellular domain of a protein involving immune response and a sequence corresponding to the DNA fragment encoding a specific region of the Fc fragment of an immunoglobulin molecule.

An recognition sequence of a restriction enzyme is introduced by PCR into the amplified immunoglobulin fusion gene and the DNA fragment having a sequence encoding a soluble extracellular domain of a protein involving the immune response. The recognition sequence is then cleaved with the restriction enzyme and the cleaved regions are ligated using ligase, thus producing a concatameric immunoglobulin fusion gene.

The immunoglobulin fusion gene may further include a signal sequence to stimulate extracellular secretion of a protein encoded thereby. For example, the CTLA-4 molecule contains a unique leader sequence having highly hydrophilic redundancy at its N-terminus, and which is abnormally long and highly water-soluble (Harper, K. et al., J. Immunol. 147:1037-1044; and Brunet, J.F. Nature 328:267-270, 1987). Generally, most

cell surface proteins or secretory proteins have a leader sequence comprising 20-24 highly hydrophobic amino acids at their N-termini. However, the CTLA-4 molecule used in the present invention comprises a total of 37 residues: 16 hydrophilic amino acids at its N-terminus, and 21 highly hydrophobic amino acids typical in its transmembrane regions.

5 In the conventional method of preparing CTLA4Ig fusion proteins, the leader sequence of the CTLA-4 molecule was substituted with a leader sequence of oncostatin M (Linsley, P.S. et al., J. Exp. Med. 174:561-569, 1991) or IL-6 (Yamada, A, et al., Microbiol. Immunol. 40:513-518, 1996). The present inventors demonstrated that a CTLA-4 molecule containing a leader sequence having a "MRTWPCTLLFFIPVFCKA" sequence

10 instead of the amino acid sequence consisting of 16 amino acids, "ACLGFRHKAQKNLAA", is preferable, and the secretion of an expressed protein to the extracellular environment is easily achieved, as disclosed in International Pat. Publication No. WO98/31820.

A recombinant expression plasmid is prepared by inserting the immunoglobulin fusion gene into a vector, and then introduced to a host cell to produce a transformant or transfectant. A concatameric fusion dimeric protein of interest may be obtained by culturing the transformant or transfectant cell and isolating and purifying a concatameric fusion protein.

15

A host cell useful for preparation of the concatameric fusion dimeric protein according to the present invention is preferably selected from among bone marrow cell lines, CHO cells, monkey COS cells, human embryonic kidney 293 cells, and baculovirus-infected insect cells. A polypeptide of interest, produced in such an expression system, is secreted to culture medium as an inclusion body. Then, the concatameric fusion dimeric protein can be purified by affinity chromatography using a protein A or protein G column. In fact, effective mammalian expression systems and such purification systems are very useful in expressing proteins involving immune response in a dimeric form, and isolation of such proteins.

20

25

Preparation of the glycosylated concatameric fusion dimeric protein according to the present invention

5 Secretory proteins produced in eukaryotic cells as host cells are modified by glycosylation. Glycosylation is known to influence in vivo stability and functionality as well as physical properties of a protein. Therefore, a preferred aspect of the present invention includes facilitating production of a concatameric fusion dimeric protein of interest using recombinant DNA techniques and the above-mentioned animal cell lines as host cells, and linking additional sugar chains to a soluble extracellular domain of a protein involving immune response.

10 Two glycosylation patterns are known. One is O-linked glycosylation, in which an oligosaccharide is linked to a serine or threonine residue, and the other is N-linked glycosylation, in which an oligosaccharide is linked to asparagine residue. N-linked glycosylation occurs at a specific amino acid sequence, particularly, Asn-X-Ser/Thr, wherein X is any amino acid excluding proline. N-linked oligosaccharide has a structure distinct from O-linked oligosaccharide, and glycosylated residues found in the N-linked type also differ from the O-linked type. For example, N-acetylgalactosamine is
15 invariably linked to serine or threonine in O-linked oligosaccharide, while N-acetylglucosamine is linked to asparagines in all of N-linked oligosaccharides. The O-linked oligosaccharides generally contain only 1-4 sugar residues. In contrast, the N-linked oligosaccharides comprise 5 or more sugar residues, essentially including N-acetylglucosamine and mannose.

20 In accordance with the present invention, to allow additional O-linked or N-linked glycosylation, one or more nucleotides in a DNA sequence encoding a soluble extracellular domain of a protein involving immune response are altered, and the resulting DNA is expressed in a suitable animal host cell to induce glycosylation using the host system. In accordance with an aspect of the present invention, the glycosylated
25 concatameric fusion dimeric protein according to the present invention may be prepared by altering a DNA sequence encoding a soluble extracellular domain of a protein involving immune response to induce or increase N-linked glycosylation by adding the sequence Asn-X-Ser/Thr.

30 Alteration of a DNA sequence to introduce glycosylation may be performed according to the conventional method common in the art. In a preferred aspect of the present invention, to protect the concatameric fusion protein, especially the two soluble

extracellular domains, from attack of intercellular proteinases and thus increase its half-life in serum, a DNA construct encoding a multiglycosylated concatameric fusion monomeric protein may be prepared using PCR, which introduces multiglycosylation sites to the joint region between two soluble extracellular domains. In a specific aspect of the present invention, glycosylation motif peptide sequences may be introduced into the concatameric fusion protein, as follows. A DNA fragment is prepared by performing PCR using a primer encoding a leader sequence of a soluble extracellular domain and EcoRI restriction site, and an antisense primer in which a portion of a nucleotide sequence encoding a portion of the 3' end of a first soluble extracellular domain and a portion of the 5' end of a second soluble extracellular domain is substituted with glycosylation motif sequences. Another DNA fragment is prepared by performing PCR using a primer in which a portion of a nucleotide sequence encoding a portion of the 3' end of a first soluble extracellular domain and a portion of the 5' end of a second soluble extracellular domain is substituted with glycosylation motif sequences, and an antisense primer encoding the 3' end of Fc portion of IgG1 and XbaI restriction site. Then, secondary PCR is carried out in a test tube using the two DNA fragments.

In accordance with an embodiment of the present invention, the soluble extracellular domains useful in the present invention include soluble extracellular domains of TNFR1, TNFR2, CD2 and CTLA-4. Their application will be described in detail with reference to accompanying figures, sequence listing and examples.

Tumor necrosis factor-alpha (TNF- α), which is known as the hormone cachectin, and tumor necrosis factor-beta (TNF- β), which is also known as lymphotoxin, are multifunctional cytokines, inducing inflammation, cellular immune response, septicemia, cytotoxicity, cachexia, rheumatoid arthritis, inflammation-related diseases (Tartaglia, L.A. et al., Immunol. Today 13:151,1992), and antiviral reaction (Butler, P., Peptide Growth Factor II, 1990, Springer-Verlag, Berlin, pp.39-70). Such actions of TNF- α and TNF- β , including cytotoxic activity, originate from their binding to TNF receptors in a trimeric form (Eck, M.J. et al., J. Biol. Chem. 267:2119, 1992). As TNF receptors, 55 kDa-type I (TNFR1 or p55) and about 75 kDa-type II (TNFR2 or p75) are known (Smith, C.A. et al., Science 248:1019, 1990; Loetscher, H. et al., Cell 61:351, 1990; and Schall et al., Cell 61:361, 1990). The two receptors have similar affinity for TNF- α and TNF- β (Schall et

al., Cell 61:361, 1990). Immunoglobulin fusion proteins of such soluble receptors have effects of inhibiting the action of TNF- α and TNF- β by inhibiting binding of TNF- α and TNF- β to their receptors on the cell surface, which is known to be effective in reducing TNF-dependent inflammation.

5 Among cell surface antigens regulating immune response, the costimulatory molecule CD2 and CTLA-4, inducing secondary stimulation to give sufficient activation of T cells, when being in a soluble form, also can be used for therapy of diverse immunological diseases according to the same method as TNF receptors. Immune
10 response is accomplished by binding of cell surface antigen molecules of antigen presenting cells (APC) to specific receptors of T lymphocytes, that is, T lymphocytes and leukocyte-function-antigen molecules of APC, and when a costimulatory signal as a secondary signal is not produced during antigen-presenting, T lymphocytes are removed by apoptosis or inhibition of clonal activation. CD2 is a leukocyte-function-antigen on
15 T lymphocytes, binding to LFA-3 on APC, and participates in adhesion and costimulation of leukocytes, as well as stimulating T cell activation through costimulation with CD28. CTLA-4 is expressed after activation of T lymphocytes, and its expression level is increased in the resting phase. CTLA-4 has a binding affinity to the B7 molecule of APC over 20 times higher than that of CD28, and transduces signals inhibiting T lymphocyte activation after binding to B7.

20 In a specific aspect of the present invention, there are provided a concatameric fusion monomeric protein TNFR1-TNFR1/Fc, designated by SEQ ID NO: 6; a concatameric fusion monomeric protein TNFR2-TNFR2/Fc, designated by SEQ ID NO: 8; a concatameric fusion monomeric protein CD2-CD2/Fc, designated by SEQ ID NO: 18; and a concatameric fusion monomeric protein CTLA4-CTLA4/Fc, designated by SEQ
25 ID NO: 20.

 In another specific aspect of the present invention, there are provided a DNA construct (TNFR1-TNFR1-IgG) encoding a concatameric fusion monomeric protein TNFR1-TNFR1/Fc, designated by SEQ ID NO: 5; a DNA construct (TNFR2-TNFR2-IgG) encoding a concatameric fusion monomeric protein TNFR2-TNFR2/Fc, designated
30 by SEQ ID NO: 7; a DNA construct (CD2-CD2-IgG) encoding a concatameric fusion monomeric protein CD2-CD2/Fc, designated by SEQ ID NO: 17; and a DNA construct

(CTLA4-CTLA4-IgG) encoding a concatameric fusion monomeric protein CTLA4-CTLA4/Fc, designated by SEQ ID NO: 19.

In a further specific aspect of the present invention, there are provided a recombinant expression plasmid pTR11Ig-Top10' operably linked to a DNA construct
5 encoding a concatameric fusion monomeric protein TNFR1-TNFR1/Fc, designated by SEQ ID NO: 5; a recombinant expression plasmid pTR22Ig-Top10' operably linked to a DNA construct encoding a concatameric fusion monomeric protein TNFR2-TNFR2/Fc, designated by SEQ ID NO: 7; a recombinant expression plasmid pCD22Ig operably linked to a DNA construct encoding a concatameric fusion monomeric protein CD2-
10 CD2/Fc, designated by SEQ ID NO: 17; and a recombinant expression plasmid pCT44Ig operably linked to a DNA construct encoding a concatameric fusion monomeric protein CTLA4-CTLA4/Fc, designated by SEQ ID NO: 19. The recombination expression plasmids are deposited in Korean Culture Center of Microorganisms (KCCM) and are assigned accession Nos. KCCM-10288, KCCM-10291, KCCM-10402 and KCCM-10400,
15 respectively. The KCCM deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

In a further specific aspect of the present invention, there are provided a mammalian host cell (e.g., TR11Ig-CHO) transformed or transfected with a recombinant
20 expression plasmid pTR11Ig-Top10' operably linked to a DNA construct encoding a concatameric fusion monomeric protein TNFR1-TNFR1/Fc, designated by SEQ ID NO: 5; a mammalian host cell (e.g., TR22Ig-CHO) transformed or transfected with a recombinant expression plasmid pTR22Ig-Top10' operably linked to a DNA construct encoding a concatameric fusion monomeric protein TNFR2-TNFR2/Fc, designated by
25 SEQ ID NO: 7; a mammalian host cell transformed or transfected with a recombinant expression plasmid pCD22Ig operably linked to a DNA construct encoding a concatameric fusion monomeric protein CD2-CD2/Fc, designated by SEQ ID NO: 17; and a mammalian host cell transformed or transfected with a recombinant expression plasmid pCT44Ig operably linked to a DNA construct encoding a concatameric fusion
30 monomeric protein CTLA4-CTLA4/Fc, designated by SEQ ID NO: 19. Chinese hamster ovary cell line TR11Ig-CHO transfected with the recombinant expression

plasmid pTR11Ig-Top10' and Chinese hamster ovary cell line TR22Ig-CHO transfected with the recombinant expression plasmid pTR22Ig-Top10' are deposited in KCCM and are assigned accession Nos. KCLRF-BP-00046 and KCLRF-BP-00049, respectively. The KCCM deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

In a still further specific aspect of the present invention, there are provided a concatameric fusion monomeric protein mgTNFR1-TNFR1/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 10; a concatameric fusion monomeric protein mgTNFR2-TNFR2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 12; a concatameric fusion monomeric protein mgCD2-CD2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 22; and a concatameric fusion monomeric protein mgCTLA4-CTLA4/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 24.

In a still further specific aspect of the present invention, there are provided a DNA construct encoding a concatameric fusion monomeric protein mgTNFR1-TNFR1/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 9; a DNA construct encoding a concatameric fusion monomeric protein mgTNFR2-TNFR2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 11; a DNA construct encoding a concatameric fusion monomeric protein mgCD2-CD2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 21; and a DNA construct encoding a concatameric fusion monomeric protein mgCTLA4-CTLA4/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 23. In order to produce a glycosylation motif peptide, a primer set (forward and reverse primers) is designed, which are complementary to a nucleotide sequence corresponding to the joint region between soluble extracellular domains of concatameric fusion proteins of TNFR/Fc, CD2/Fc and CTLA4/Fc, as well as containing codons encoding asparagine (N) (ATT and AAC) or codons encoding serine (S) and threonine (T) (TCC; and ACC, ACG and ACA, respectively), with which any codon in the concatameric fusion protein gene may be substituted. When designing the primer, selection of one among a plurality of amino acid sequences may be determined

depending on a condition allowing minimum substitution of the nucleotide sequence and melting temperature (T_m) of each primer.

In a still further specific aspect of the present invention, there are provided a recombinant expression plasmid pTR11Ig-MG operably linked to a DNA construct
5 encoding a concatameric fusion monomeric protein mgTNFR1-TNFR1/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 9; a recombinant expression plasmid pTR22Ig-MG operably linked to a DNA construct encoding a concatameric fusion monomeric protein mgTNFR2-TNFR2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 11; a recombinant expression plasmid pCD22Ig-MG operably
10 linked to a DNA construct encoding a concatameric fusion monomeric protein mgCD2-CD2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 21; and a recombinant expression plasmid Pct44Ig-MG operably linked to a DNA construct encoding a concatameric fusion monomeric protein mgCTLA4-CTLA4/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 23. The recombination
15 expression plasmids are deposited in Korean Culture Center of Microorganisms (KCCM) and are assigned accession Nos. KCCM-10404, KCCM-10407, KCCM-10401 and KCCM-10399, respectively. The KCCM deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

In a still further specific aspect of the present invention, there are provided a
20 mammalian host cell transformed or transfected with a recombinant expression plasmid pTR11Ig-MG operably linked to a DNA construct encoding a concatameric fusion monomeric protein mgTNFR1-TNFR1/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 9; a mammalian host cell transformed or transfected with a
25 recombinant expression plasmid pTR22Ig-MG operably linked to a DNA construct encoding a concatameric fusion monomeric protein mgTNFR2-TNFR2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 11; a mammalian host cell transformed or transfected with a recombinant expression plasmid pCD22Ig-MG operably
30 linked to a DNA construct encoding a concatameric fusion monomeric protein mgCD2-CD2/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 21; and a mammalian host cell transformed or transfected with a recombinant expression plasmid

Pct44Ig-MG operably linked to a DNA construct encoding a concatameric fusion monomeric protein mgCTLA4-CTLA4/Fc containing glycosylation motif peptides, designated by SEQ ID NO: 23.

5 The concatameric fusion dimeric proteins of the present invention may be isolated from culture medium after culturing the transformants or transfectants according to the present invention. The concatameric fusion dimeric proteins may participate in immune response, as described in Table 1, above, and are thus useful as therapeutic agents, diagnostic agents and laboratory tools according to the kinds of the protein, and their use is well known to those of ordinary skill in the art. In particular, when being
10 used as therapeutic agents, the concatameric fusion dimeric proteins may be applied at an therapeutically effective amount common in the art, and it will be understood that such an amount may vary depending on diverse factors including activity of the used compound, patient's age, body weight, health state, sex and diet, administration time, administration route, combination of drugs, and pathogenic state of a specific disease to be prevented or
15 treated. In addition, when being used as therapeutic agents, it will be understood that the concatameric fusion dimeric proteins according to the present invention may be applied by the typical methods and routes for administration of proteins involving immune response, which are known to those of ordinary skill in the art.

20 The present invention will be explained in more detail with reference to the following examples in conjunction with the accompanying drawings. However, the following examples are provided only to illustrate the present invention, and the present invention is not limited to them. For convenience in describing the present invention, information on DNA constructs, recombinant expression plasmids and transformed cell lines, which are prepared according to the Examples, below, and the used primers and
25 accession numbers is summarized in Tables 8 and 9, below.

TABLE 8
Information on DNA constructs and accession Nos.

DNA construct name	SEQ ID No.		Deposition of genes		Deposition of cell lines	
	DNA	Protein	Designation	Accession No.	Designation	Accession No.
TNFR1-IgG	1	2				
TNFR2-IgG	3	4				
TNFR1-TNFR1-IgG	5	6	pTR11Ig-Top10'	KCCM 10288	TR11Ig-CHO	KCLRF-BP-00046
TNFR2-TNFR2-IgG	7	8	pTR22Ig-Top10'	KCCM 10291	TR22Ig-CHO	KCLRF-BP-00049
mgTNFR1-TNFR1-IgG	9	10	pTR11Ig-MG	KCCM 10404		
mgTNFR2-TNFR2-IgG	11	12	pTR22Ig-MG	KCCM 10407		
CD2-IgG	13	14				
CTLA4-IgG	15	16				
CD2-CD2-IgG	17	18	pCD22Ig	KCCM 10402		
CTLA4-CTLA4-IgG	19	20	pCT44Ig	KCCM 10400		
mgCD2-CD2-IgG	21	22	pCD22Ig-MG	KCCM 10401		
mgCTLA4-CTLA4-IgG	23	24	pCT44Ig-MG	KCCM 10399		

TABLE 9
Information for primers

Primer name	SEQ ID No.	Description
Oligo TNFR-EDF-EcoRI	25	Containing 5' end of the extracellular domain of TNFR1 and an EcoRI site
Oligo TNFR-EDR-IgGh	26	Reverse primer containing 3' end of the extracellular domain of TNFR1 and the hinge region of IgG
Oligo IgG1-T1F	27	Containing 5' end of the hinge region of IgG and 3' end of TNFR1
Oligo IgG1-R-XbaI	28	Reverse primer containing 3' end of the hinge region of IgG and a XbaI site
Oligo TNFR2-EDF-EcoRI	29	Containing 5' end of the extracellular domain of TNFR2 and an EcoRI site
Oligo TNFR2-EDR-IgGh	30	Reverse primer containing 3' end of the extracellular domain of TNFR2 and the hinge region of IgG
Oligo IgG1-T2F	31	Containing 5' end of the hinge region of IgG and 3' end of TNFR2
Oligo TNFR1-CF-BamHI	32	Containing 5' end of the extracellular domain of TNFR1 and a BamHI site; and used for preparation of a concatamer
Oligo TNFR1-NR-BamHI	33	Reverse primer containing 3' end of the extracellular domain of TNFR1 and a BamHI site; and used for preparation of a concatamer
Oligo TNFR2-CF-BamHI	34	Containing 5' end of the extracellular domain of TNFR2 and a BamHI site; and used for preparation of a concatamer
Oligo TNFR2-NR-BamHI	35	Reverse primer containing 3' end of the extracellular domain of TNFR2 and a BamHI site; and used for preparation of a concatamer
Oligo mgTNFR1-TNFR1-IgG-F	36	Primer for mutagenesis, containing a sequence capable of inserting glycosylation sites into the joint region of TNFR1-TNFR1, and sequences corresponding to 3' end and 5' end of TNFR1; and used for preparation of a MG (multiglycosylation) form
Oligo mgTNFR1-TNFR1-IgG-R	37	Reverse primer for mutagenesis, containing a sequence capable of inserting glycosylation sites into the joint region of TNFR1-TNFR1, and sequences corresponding to 3' end and 5' end of TNFR1; and used for preparation of a MG form
Oligo mgTNFR2-TNFR2-IgG-F	38	Primer for mutagenesis, containing a sequence capable of inserting glycosylation sites into the joint region of TNFR2-TNFR2, and sequences corresponding to 3' end and 5' end of TNFR2; and used for preparation of a MG form
Oligo mgTNFR2-TNFR2-IgG-R	39	Reverse primer for mutation, containing a sequence capable of inserting glycosylation sites into the joint region of TNFR2-TNFR2, and sequences corresponding to 3' end and 5' end of TNFR2; and used for preparation of a MG form
Oligo CD2F-EcoRI	40	Containing 5' end of the extracellular domain of CD2 and a EcoRI site
Oligo CD2R-RstI	41	Containing 3' end of the extracellular domain of CD2 and a PstI site
Oligo IgG-F-PstI	42	Containing 5' end of the hinge region of IgG and a PstI site
Oligo CTLA4F-EcoRI	43	Containing 5' end of the extracellular domain of CTLA-4 and a EcoRI site
Oligo CTLA4R-PstI	44	Containing 3' end of the extracellular domain of CTLA-4 and a PstI site
Oligo CD2-NT-F	45	Containing 5' end of the extracellular domain of CD2; and used for preparation of a concatamer
Oligo CD2-CT-R	46	Reverse primer containing 3' end of the extracellular domain of CD2; and used for preparation of a concatamer
Oligo CTLA4-NT-F	47	Containing 5' end of the extracellular domain of CTLA-4; and used for preparation of a concatamer
Oligo CTLA4-CT-R	48	Reverse primer containing 3' end of the extracellular domain of CTLA-4; and used for preparation of a concatamer
Oligo mgCD2-CD2-IgG-F	49	Used for preparation of a MG (multiglycosylation) form of CD2-CD2-IgG
Oligo mgCD2-CD2-IgG-R	50	Reverse primer used for preparation of a MG (multiglycosylation) form of CD2-CD2-IgG
Oligo mgCTLA4-CTLA4-IgG-F	51	Used for preparation of a MG (multiglycosylation) form of CTLA4-CTLA4-IgG
Oligo mgCTLA4-CTLA4-IgG-R	52	Reverse primer used for preparation of a MG (multiglycosylation) form of CTLA4-CTLA4-IgG

EXAMPLE 1**Human TNFR**

5 A. Manufacture of a DNA construct encoding simple fusion monomeric protein of TNFR1/Fc (Fig. 1 and Fig. 5)

a. DNA fragment encoding soluble extracellular domain of TNFR1

10 A fusion gene encoding soluble extracellular domain of type I human TNF receptor (TNFR1, p55) and Fc fragment of human immunoglobulin G1 was constructed by the Polymerase Chain Reaction (PCR) method described in the prior art (Holten et al., Biotechniques 8:528, 1990).

15 A DNA fragment encoding soluble extracellular domain of TNFR1 was constructed by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 25) with EcoRI restriction site and the sequence encoding leader sequence (the sequence of amino acids 1-20 of SEQ ID NO: 2), and an antisense primer (the sequence of nucleotide of SEQ ID NO: 26) with the sequence encoding a part of 3' ends of the said soluble extracellular domain of TNFR1 (TNFR1-ED) and 5' ends of the hinge region of immunoglobulin G1 (IgG1). The template cDNA for this reaction was constructed by reverse transcription PCR (RT-PCR) of mRNA extracted from monocyte (T lymphocyte) of healthy adults.

20 After blood of healthy adults was extracted and diluted to 1:1 with RPMI-1640 (Gibco BRL, USA), the layer of T lymphocyte which formed at upper part was obtained by density gradient centrifugation using Ficoll-hypaque (Amersham, USA). In order to make the concentration of the cell to 5×10^5 cells/ml, the cell was washed with RPMI-1640 for 3 times, and RPMI-1640 culture media containing 10% Fetal Bovine Serum (FBS, Gibco BRL, USA) was added, then cultured at 37°C for two days in the 5% CO₂ incubator after
25 adding leukoagglutinin to 3.5ug/ml (Pharmacia, USA).

The mRNAs were purified using Tri-Reagent (MRC, USA) mRNA purification kit. First, 2×10^7 of human T lymphocyte was washed with Phosphate Buffered Saline (PBS, pH7.2) for 3 times, and then 1ml of Tri-Reagent was mixed for several times to dissolve RNA. After adding 0.2ml of chloroform to this tube and mixing thoroughly, this tube was incubated at room temperature (RT) for 15 min, then centrifuged at 15,000 rpm, 4°C for 15 min. The upper part of the solution was transferred to a 1.5ml tube, and 0.5ml of isopropanol was added, and then centrifuged at 15,000 rpm, 4°C for 15 min. After the supernatant was discarded, the pellet was resuspended with 1ml of 3° distilled water treated with 75% ethanol-25% DEPC (Sigma, USA), and then centrifuged at 15,000 rpm, 4°C for 15 min. After the supernatant was removed completely and dried in the air to remove ethanol residue, RNA was resuspended with 50µl of 3° distilled water treated with DEPC.

The primary cDNA was synthesized by mixing 2µg of purified mRNA and 1µl of oligo dT (dT30, Promega, USA) primer to 10µM in 1.5ml tube, heating at 70°C for 2 min, and cooling in ice for 2 min. After that, this mixture was added with 200U of M-MLV reverse transcriptase (Promega, USA), 10µl of 5 x reaction buffer (250mM Tris-HCl, pH 8.3, 375mM KCl, 15mM MgCl₂, and 50mM DTT), 1µl of dNTP (10mM each, Takara, Japan), and DEPC-treated 3° distilled water to 50µl, then reacted at 42°C for 1 hour.

b. DNA fragment encoding Fc fragment of immunoglobulin

A DNA fragment encoding Fc fragment of immunoglobulin G1 was constructed by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 27) with the sequence encoding a part of 3' ends of the said soluble extracellular domain of TNFR and 5' end of the hinge region of immunoglobulin G1 (IgG1), and an antisense primer (the sequence of nucleotide of SEQ ID NO: 28) with XbaI restriction site and the sequence encoding 3' ends of IgG1 Fc. The template cDNA for this reaction was constructed by RT-PCR of mRNA extracted from peripheral blood cell (B lymphocyte) of convalescent patients with pyrexia of unknown origin.

c. DNA construct encoding simple fusion monomeric protein of TNFR1/Fc

After DNA fragment encoding soluble extracellular domain of TNFR1 and DNA fragment encoding Fc fragment of immunoglobulin produced as described above were mixed in the same tube, complementary binding between the common sequence (the sequence including 3' end of soluble extracellular domain of TNFR1 and 5' end of IgG1 hinge region) was induced. Using this mixture as a template, DNA construct including DNA fragment encoding soluble extracellular domain of TNFR1 and DNA fragment encoding IgG1 Fc fragment was amplified by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 25) with the sequence encoding 5' end of TNFR1 and another primer (the sequence of nucleotide of SEQ ID NO: 28) with the sequence encoding 3' end of IgG1 Fc. The constructed gene included a leader sequence to facilitate secretion of protein after expression.

d. Cloning of the DNA construct encoding simple fusion monomeric protein of TNFR1/Fc

DNA construct encoding simple fusion monomeric protein of TNFR1/Fc as described above was restricted with EcoRI and XbaI, and cloned by inserting into a commercially available cloning vector, pBluescript KS II (+) (Stratagene, USA), at EcoRI/XbaI site. The sequence of a total coding region was identified by DNA sequencing (SEQ ID NO: 1). This produced fusion protein was designated TNFR1/Fc as simple fusion monomeric protein, and the elliptical shape shown in Figure 1 represents the structure of a primary expression product of the fusion gene. The deduced amino acid sequence of simple fusion monomeric of TNFR1/Fc corresponded to SEQ ID NO: 2.

B. Manufacture of a DNA construct encoding simple fusion monomeric protein of TNFR2/Fc (Fig. 1 and Fig. 5)

a. DNA fragment encoding soluble extracellular domain of TNFR2

5 A fusion gene encoding soluble extracellular domain of type II human TNF receptor (TNFR2, p75) and Fc fragment of human immunoglobulin G1 was constructed by the same method as that of TNFR1/Fc.

10 A DNA fragment encoding soluble extracellular domain of TNFR2 was constructed by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 29) with EcoRI restriction site and the sequence encoding leader sequence (the sequence of amino acids 1-22 of SEQ ID NO: 4), and an antisense primer (the sequence of nucleotide of SEQ ID NO: 30) with the sequence encoding a part of 3' ends of said soluble extracellular domain of TNFR2 (TNFR2-ED) and 5' ends of the hinge region of immunoglobulin G1 (IgG1). The template cDNA for this reaction was constructed by RT-PCR of mRNA extracted from monocyte (T lymphocyte) of healthy adults.

15 b. DNA construct encoding simple fusion monomeric protein of TNFR2/Fc

20 After DNA fragment encoding soluble extracellular domain of TNFR2 and DNA fragment encoding Fc fragment of immunoglobulin G1 produced as described above were mixed in the same tube, complementary binding between the common sequence (the sequence including 3' end of soluble extracellular domain of TNFR2 and 5' end of IgG1 hinge region) was induced. Using this mixture as a template, DNA construct including DNA fragment encoding soluble extracellular domain of TNFR2 and encoding and DNA fragment encoding IgG1 Fc fragment was amplified by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 29) with the sequence encoding 5' end of TNFR2 and another primer (the sequence of nucleotide of SEQ ID NO: 28) with the sequence encoding 3' end of IgG1 Fc. The constructed gene includes a sequence to facilitate secretion of protein after expression.

25

c. Cloning of the DNA construct encoding simple fusion monomeric protein of TNFR2/Fc

DNA construct encoding simple fusion monomeric protein of TNFR2/Fc as described above was restricted with EcoRI and XbaI, and cloned by inserting into a commercially available cloning vector, pBluescript KS II (+) (Stratagene, USA), at EcoRI/XbaI site. The sequence of a total coding region was identified by DNA sequencing (SEQ ID NO: 3). This produced fusion protein was designated TNFR2/Fc as simple fusion monomeric protein, and the elliptical shape shown in Figure 1 represents the structure of a primary expression product of the fusion gene. The deduced amino acid sequence of simple fusion monomeric of TNFR2/Fc corresponded to SEQ ID NO: 4.

C. Manufacture of a DNA construct encoding concatameric fusion monomeric protein of TNFR1-TNFR1/Fc (Fig. 2 and Fig. 5)

In order to manufacture a fusion gene comprising the concatameric shape in soluble extracellular domain of TNFR1, i.e. the DNA construct encoding concatameric fusion monomeric protein of TNFR1-TNFR1/Fc, BamHI restriction site was inserted respectively into the sequence of soluble extracellular domain of TNFR1 and DNA construct as produced as above encoding simple fusion monomeric protein of TNFR1/Fc by PCR, and then regions of each fragments restricted by BamHI were linked by ligase. The DNA construct, encoding simple fusion monomeric protein of TNFR1/Fc produced as above, was used as the template of this reaction.

The fragment of the soluble extracellular domain of TNFR1 with BamHI restriction site at 3' end was amplified by PCR using a primer corresponding to the nucleotide of SEQ ID NO: 25 and another primer corresponding to the nucleotide sequence of SEQ ID NO: 33, and the other fragment of simple fusion monomeric protein of TNFR1/Fc with BamHI restriction site at 5' end was amplified by PCR using a primer

corresponding to the nucleotide of SEQ ID NO: 28 and another primer corresponding to the nucleotide sequence of SEQ ID NO: 32, respectively. PCR was performed by adding 1 μ l of primary cDNA, 2U of Pfu DNA polymerase (Stratagene, USA), 10 μ l of 10X reaction buffer [200mM Tris-HCl, pH 8.75, 100mM (NH₄)₂SO₄, 100mM KCl, 20mM MgCl₂], 1% TritonTM X-100, 1mg/ml BSA, 3 μ l primer 1 (10 μ M), 3 μ l primer 2 (10 μ M), 2 μ l dNTP (10mM each), and 3° distilled water to 100 μ l. The reaction condition was as follows; 94°C, 5 min; 95°C, 1 min; 58°C, 1 min 30 sec; 72°C, 1 min for 31 cycles; and 72°C, 15 min to make PCR product with complete blunt end.

After electrophorized on 0.8% agarose gel, the PCR product was purified by Qiaex II gel extraction kit (Qiagen, USA). The purified PCR product was restricted by BamHI and extracted by phenol-chloroform extraction methods. Subsequently, two kinds of DNA fragments restricted by BamHI were linked by ligase.

D. Manufacture of a DNA construct encoding concatameric fusion monomeric protein of TNFR2-TNFR2/Fc (Fig. 2 and Fig. 5)

After a BamHI restriction site was inserted respectively into the sequence of the soluble extracellular domain of TNFR21 and the DNA construct produced as described above encoding simple fusion monomeric protein of TNFR2/Fc by PCR, a DNA construct encoding concatameric fusion monomeric protein of TNFR2-TNFR2/Fc was manufactured by linking the regions of each fragments restricted by BamHI by ligase.

A fragment of soluble extracellular domain of TNFR2 with BamHI restriction site at 3' end was amplified using a primer corresponding the sequence of SEQ ID NO: 34 and SEQ ID NO: 35. PCR was performed as that of TNFR1, except that a DNA construct encoding simple fusion monomeric protein of SEQ ID NO: 3 produced as above was used as a template. The PCR product was purified by the method as that of TNFR1.

E. DNA construct encoding concatameric fusion monomeric protein of TNFR1-TNFR1/Fc with glycosylation motif.

5 A DNA fragment was manufactured by PCR using an antisense primer (the sequence of nucleotide of SEQ ID NO: 37) with the sequence encoding the part (the sequence of nucleotide 565-591 of SEQ ID NO: 5) of 3' end of the first soluble extracellular domain of TNFR1, except the sequence of hydrophobic peptide region (the sequence of amino acid 197-216 of SEQ ID NO: 6) at the junction of soluble extracellular domain of TNFR1 and the part (the sequence of nucleotide 649-681 of SEQ ID NO: 5) of 5' end of the
10 second soluble extracellular domain of TNFR1, and another primer (the sequence of nucleotide of SEQ ID NO: 25) with the sequence encoding EcoRI restriction site and leader sequence.

In addition, the total four amino acid sequences encoding glycosylation site (the sequence of amino acids 189-191, 192-194, 198-200, and 204-206 of SEQ ID NO: 10)
15 were inserted by manufacturing the primer as above (the sequence of nucleotide of SEQ ID NO: 36 and 37) corresponding the substitution of the nucleotide 565-567 (CTG, Leu), 574-576 (ACG, Thr), 652-654 (CTA, Leu), and 670-672 (AGA, Arg) of SEQ ID NO: 5 with the nucleotide of AAC (Asn, N); the nucleotide of 571-573 (TGC, Cys) and 580-582 (TTG, Leu) of SEQ ID NO: 5 with the nucleotide of ACC (Thr, T); the nucleotide of 658-660
20 (GAC, Asp) with the nucleotide of TCC (Ser, S).

In this reaction, the gene (the nucleotide of SEQ ID NO: 5) encoding concatameric shape of TNFR1-TNFR1/Fc was used as a template. During the primary PCR, only the half of the antisense primer was induced to bind the gene encoding concatameric shape of TNFR1-TNFR1/Fc used as a template, and, as chain reaction was
25 proceeding, the unbound part to the template was induced to form a complete double-stranded DNA by polymerase, and then this was capable of producing the DNA fragment with state of linkage of the sequence of 5' end encoding the part of the second soluble

extracellular domain and the sequence of 3' end encoding TNFR1 extracellular domain including leader sequence. Therefore, a part of the sequence of 5' end encoding the second soluble extracellular domain has the function that was capable of binding to the second DNA fragment as follows.

5 The second DNA fragment was manufactured by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 36) with the sequence encoding the part (the sequence of nucleotide 565-591 of SEQ ID NO: 5) of 3' end of the first soluble extracellular domain of TNFR1 and the part (the sequence of nucleotide 649-681 of SEQ ID NO: 5) of 5' end of the second soluble extracellular domain of TNFR1, and an antisense primer (the
10 sequence of nucleotide of SEQ ID NO: 28) with the sequence encoding a XbaI restriction site and 3' end of IgG1 Fc. This reaction was also performed as described above, that is, only the half of antisense primer was induced to bind the template, and consequently, DNA fragment like that described above had the sequence encoding 5' end of TNFR1 extracellular including the part of 3' end of the first soluble extracellular domain.

15 Subsequently, resulting from two kinds of DNA fragments as PCR described as above were mixed in the same tube, induced to bind between common sequences, and fused by PCR using primers (the sequence of nucleotide of SEQ ID NO: 25 and 28) encoding 5' and 3' end of each concatameric genes, and the product was designated mgTNFR1-TNFR1-IgG.

20

F. DNA construct encoding concatameric fusion monomeric protein of TNFR2-TNFR2/Fc with glycosylation motif.

 A DNA fragment was manufactured by PCR using an antisense primer (the
25 sequence of nucleotide of SEQ ID NO: 39) with the sequence encoding the part (the sequence of nucleotide 586-606 of SEQ ID NO: 7) of 3' end of first soluble extracellular domain of TNFR2, except the sequence of hydrophobic peptide region (the sequence of

amino acid 203-263 of SEQ ID NO: 8) at the junction of soluble extracellular domain of TNFR2 and the part (the sequence of nucleotide 790-807 of SEQ ID NO: 7) of 5' end of second soluble extracellular domain of TNFR2, and another primer (the sequence of nucleotide of SEQ ID NO: 29) with the sequence encoding EcoRI restriction site and leader sequence.

In addition, the total two amino acid sequences encoding glycosylation site (the sequence of amino acids 199-201 and 206-208 of SEQ ID NO: 12) were inserted by manufacturing the primer as described above (the sequence of nucleotide of SEQ ID NO: 38 and 39) corresponding to the substitution of the nucleotide 595-597 (GTC, Val) and 799-801 (GGG, Gly) SEQ ID NO: 7 with the nucleotide of AAC (Asn, N).

In this reaction, the gene (the nucleotide of SEQ ID NO: 7) encoding concatameric shape of TNFR2-TNFR2/Fc was used as a template. During the primary PCR, only the half of antisense primer was induced to bind the gene encoding concatameric shape of TNFR2-TNFR2/Fc used as a template, and, as the chain reaction was proceeding, the unbound part to the template was induced to form a complete double-stranded DNA by polymerase, and thus this was capable of producing the DNA fragment with a state of linkage of the sequence of 5' end encoding the part of the second soluble extracellular domain and the sequence of 3' end encoding TNFR2 extracellular domain including the leader sequence. Therefore, a part of the sequence of 5' end encoding the second soluble extracellular domain has the function that was capable of binding to the second DNA fragment as follows.

The second DNA fragment was manufactured by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 38) with the sequence encoding the part (the sequence of nucleotide 586-606 of SEQ ID NO: 7) of 3' end of the first soluble extracellular domain of TNFR2 and the part (the sequence of nucleotide 790-807 of SEQ ID NO: 7) of 5' end of the second soluble extracellular domain of TNFR2, and an antisense primer (the sequence of nucleotide of SEQ ID NO: 28) with the sequence encoding a XbaI restriction

site and 3' end of IgG1 Fc. This reaction was also performed, that is, only the half of antisense primer was induced to bind the template, and consequently, DNA fragment like that described above had the sequence encoding 5' end of TNFR2 extracellular including the part of 3' end of first soluble extracellular domain.

5 Subsequently, resulting from two kinds of DNA fragments as PCR produced as above were mixed in the same tube, induced to bind between common sequences, and fused by PCR using primers (the sequence of nucleotide of SEQ ID NO: 29 and 28) encoding 5' and 3' end of each concatameric genes, and the product was designated mgTNFR2-TNFR2-IgG.

10 G. Cloning of DNA constructs encoding concatameric fusion monomeric protein of TNFR-TNFR/Fc and their glycosylated forms

DNA constructs encoding concatameric fusion monomeric protein of TNFR-TNFR/Fc and their glycosylated forms as above were cloned by inserting into pBluescript
15 KS II (+) (Stratagene, USA) at EcoRI/XbaI site. These produced fusion proteins were designated TNFR1-TNFR1/Fc and TNFR2-TNFR2/Fc as concatameric fusion monomeric protein, and designated mgTNFR1-TNFR1/Fc and mgTNFR2-TNFR2/Fc as their glycosylated forms. The deduced amino acid sequences corresponded to SEQ ID NO: 6, 8,
20 10, and 12.

After 10 μ g of pBluescript KS II (+) (Stratagene, USA) used as a vector was mixed with 15U of EcoRI, 15U of XbaI, 5 μ l of 10X reaction buffer (100mM Tris-HCl, pH 7.5, 100mM MgCl₂, 10mM DTT, 500mM NaCl), 5 μ l of 0.1% BSA (Takara, Japan), and 3 $^\circ$ distilled water to 50 μ l, DNA was restricted by incubation at 37 $^\circ$ C for 2 hrs. After
25 electrophorized on 0.8% agarose gel, the PCR product was purified by Qiaex II gel extraction kit (Qiagen, USA).

After 100ng of pBluescript KS II (+) (Stratagene, USA) restricted by EcoRI and XbaI was mixed with 20ng of PCR product restricted by the restriction enzyme, 0.5U of T4 DNA ligase (Amersham, USA), 1µl of 10X reaction buffer (300mM Tris-HCl, pH 7.8, 100mM MgCl₂, 100mM DTT, 10mM ATP) and 3° distilled water were added to 10µl, and the mixture was incubated in the water bath at 16°C for 16 hrs. E. coli Top10 (Novex, USA) was made to competent cell by the method of rubidium chloride (RbCl, Sigma, USA) and transformed, then spread on the solid LB media including 50µg/ml of ampicillin (Sigma, USA) and incubated at 37°C for 16 hrs. Formed colonies were inoculated in 4ml of liquid LB media including 50µg/ml of ampicillin and incubated at 37°C for 16 hrs. Plasmid was purified by the method of alkaline lysis according to Sambrook et al. (Molecular cloning, Cold Spring Harbor Laboratory press, p1.25-1.31, p1.63-1.69, p7.26-7.29, 1989) from 1.5ml of that, and the existence of cloning was confirmed by the restriction of EcoRI and XbaI.

The sequence of a total coding region was identified by the DNA sequencing method of dideoxy chain termination method (Sanger et al., Proc. Natl. Acad. Sci., 74:5483, 1977) as follows. The DNA sequencing reaction was performed according to the manual using a plasmid purified by alkaline lysis method as described above and SequenaseTM ver 2.0 (Amersham, USA). After the reaction mixture as above was loaded on 6% polyacrylamide gel and electrophorized for 2 hrs at constant voltage of 1,800~2,000 V and 50°C, DNA sequence was identified by exposing to X-ray film (Kodak, USA) after the gel was dried out.

EXAMPLE 2 AND 3

CD2 and CTLA4

DNA fragments encoding soluble extracellular domain of CD2 and CTLA4 were constructed by PCR using a primer [CD2(the sequence of nucleotide of SEQ ID NO:

40), and CTLA4(the sequence of nucleotide of SEQ ID NO: 43)] with EcoRI restriction site and the coding sequence [CD2 (the sequence of nucleotide of SEQ ID NO: 13), and CTLA4 (the sequence of nucleotide of SEQ ID NO: 15)] encoding the leader sequence [CD2(the sequence of amino acid 1-24 of SEQ ID NO: 14), and CTLA4(the sequence of amino acid 1-21 of SEQ ID NO: 16)], and an antisense primer [CD2(the sequence of nucleotide of SEQ ID NO: 41), and CTLA4(the sequence of nucleotide of SEQ ID NO: 44)] with PstI restriction site and the sequence [CD2(the sequence of nucleotide of SEQ ID NO: 13), and CTLA4(the sequence of nucleotide of SEQ ID NO: 15)] encoding 3' end of the soluble extracellular domain of the proteins as described above. The template cDNA for this reaction was constructed by reverse transcription PCR (RT-PCR) of mRNA extracted from the monocyte (T lymphocyte) of healthy adults.

Also, a DNA fragment encoding Fc fragment of immunoglobulin G1 was constructed by PCR using a primer (the sequence of nucleotide of SEQ ID NO: 42) with PstI restriction site and the sequence encoding 5' ends of constant region of IgG1, and an antisense primer (the sequence of nucleotide of SEQ ID NO: 28) with XbaI restriction site and the sequence encoding 3' ends of IgG1 Fc. The template cDNA for this reaction was constructed by RT-PCR of mRNA extracted from peripheral blood cell (B lymphocyte) of convalescent patients with unknown fever.

Subsequently, both DNA fragment encoding soluble extracellular domain of CD2 and CTLA4 and DNA fragment encoding Fc fragment of immunoglobulin G1 produced as described above were restricted by PstI, and then the simple dimeric shape of CD2/Fc and CTLA4/Fc genes were constructed by linkages using T4 DNA ligase. The constructed genes included a leader sequence to facilitate secretion of protein after expression.

DNA constructs as described above were restricted by restriction enzyme of EcoRI and XbaI, and cloned by inserting into a commercially available cloning vector, pBluescript KS II (+) (Stratagene, USA) at EcoRI/XbaI site. The sequence of a total coding

region was identified by DNA sequencing (SEQ ID NO: 13 and 15). These produced fusion proteins were designated CD2/Fc and CTLA4/Fc, and the deduced amino acid sequences of these corresponded to SEQ ID NO: 14 and 16.

5 PCR was performed by adding 1 μ l of primary cDNA, 2U of Pfu DNA polymerase (Stratagene, USA), 10 μ l of 10X reaction buffer [200mM Tris-HCl, pH 8.75, 100mM (NH₄)₂SO₄, 100mM KCl, 20mM MgCl₂], 1% TritonTM X-100, 1mg/ml BSA, 3 μ l primer 1 (10 μ M), 3 μ l primer 2 (10 μ M), 2 μ l dNTP (10mM each), and 3° distilled water to 100 μ l. The reaction condition was as follows; 94°C, 5 min; 95°C, 1 min; 58°C, 1 min 30 sec; 72°C, 1 min for 31 cycles; and 72°C, 15 min to make PCR product with complete blunt
10 end.

The fusion genes with concatameric shape of CD2-CD2/Fc and CTLA4-CTLA4/Fc were constructed as follows.

In order to manufacture fusion gene comprising the concatameric shape in soluble extracellular domain of CD2 and CTLA4, the sequences of soluble extracellular
15 domain of CD2 and CTLA4 were inserted by blunt-end ligation using ligase at the junction between extracellular domain and immunoglobulin of fusion genes in the shape of simple dimer with blunt end, using PstI restriction enzyme and T4 DNA polymerase. Specifically, DNA constructs were constructed by PCR using a primer [CD2(the sequence of nucleotide of SEQ ID NO: 13) and CTLA4(the sequence of nucleotide of SEQ ID NO: 48)] with the
20 coding sequence [CD2(the sequence of nucleotide of SEQ ID NO: 13) and CTLA4(the sequence of nucleotide of SEQ ID NO: 15)] encoding the end of leader sequence [CD2(the sequence of amino acid 25 of SEQ ID NO: 14) and CTLA4(the sequence of amino acid 22 of SEQ ID NO: 16)] of soluble extracellular domain, and an antisense primer [CD2(SEQ ID NO: 46) and CTLA4(SEQ ID NO: 48)] with the sequence [CD2(the sequence of nucleotide
25 of SEQ ID NO: 13) and CTLA4(the sequence of nucleotide of SEQ ID NO: 15)] encoding 3' end of soluble extracellular domain as above. The simple fusion monomeric genes [CD2/Fc (the sequence of nucleotide of SEQ ID NO: 13) and CTLA4/Fc (the sequence of

nucleotide of SEQ ID NO: 15)] described as above were used as the template of this reaction.

Also, CD2/Fc and CTLA4/Fc, which were inserted in pBluescript KS II (+) in the shape of simple monomeric form, were made to have 3' overhang end using the restriction enzyme of PstI. The cut end of 3' overhang was partially deleted to form a blunt end by treating T4 DNA polymerase. In order to manufacture fusion genes in the shape of concatamer in soluble extracellular domain, the soluble extracellular domains of CD2 and CTLA4 produced by PCR as described above were cloned by inserting into cut ends of simple monomeric gene made as blunt end. These produced fusion proteins were designated CD2-CD2/Fc and CTLA4-CTLA4/Fc as concatameric fusion monomeric protein, and their deduced amino acid sequences corresponded SEQ ID NO: 18 and 20, respectively.

The concatameric fusion genes in the shape of multiglycosylated form were constructed as follows.

The glycosylation motif was inserted by secondary PCR with mixing in the same tube of a DNA fragment produced by PCR using a primer including EcoRI restriction site and the soluble extracellular domain with leader sequence, and an antisense primer with the sequence encoding the part of 3' end of the first soluble extracellular domain of concatameric shape of fusion gene and the part of 5' end of the second soluble extracellular domain with the nucleotide of substituted glycosylation motif; and other DNA fragment produced by PCR using a primer with the sequence encoding the part of 3' end of the first soluble extracellular domain of concatameric shape of fusion gene and the part of 5' end of the second soluble extracellular domain with the nucleotide of substituted glycosylation motif, and an antisense primer with the sequence encoding 3' end of Fc fragment of immunoglobulin G1 and XbaI restriction site.

In the case of concatameric fusion gene of CD2/Fc and CTLA4/Fc, the glycosylation motif was inserted by PCR using modified primers as the same methods as

that of TNFR/Fc described as above, but it was different from the case of TNFR/Fc that the amino acid sequence of binding to soluble extracellular domain of CD2 and CTLA4 was retained as the same.

In the process of multiglycosylation of the concatameric fusion protein of CD2/Fc and CTLA4/Fc, the case of CD2/Fc was completed by inserting the total two glycosylation motif peptide region (the sequence of amino acid of 200-202 and 206-208 of SEQ ID NO: 22) using a manufactured primer including the substitution of the nucleotide of 598-600 (CCT, Pro) and 616-618 (GAG, Glu) of SEQ ID NO: 17 with AAT (Asn, N), and the case of CTLA4/Fc was completed by inserting the total three glycosylation motif peptide region (the sequence of amino acid of 136-138, 142-144, and 147-149 of SEQ ID NO: 24) using a manufactured primer (SEQ ID NO: 51 and 52) including the substitution of the nucleotide of 403-405 (GTA, Val) and 424-426 (CCA, Pro) of SEQ ID NO: 19 with AAT (Asn, N); the nucleotide of 409-411 (GAT, Asp) and 445-447 (GTG, Val) with ACA (Thr, T) and ACG (Thr, T), respectively. These produced fusion proteins were designated mgCD2-CD2/Fc and mgCTLA4-CTLA4/Fc as concatameric fusion monomeric protein, and their deduced amino acid sequences corresponded to SEQ ID NO: 22 and 24, respectively.

EXAMPLE 4

Expression and purification of simple/concatameric fusion dimeric protein of TNFR/Fc

In order to express the fusion proteins in CHO-K1 cell (ATCC CCL-61, Ovary, Chinese hamster, *Cricetulus griseus*), after pBluescript KS II (+) plasmid DNA including TNFR/Fc fusion gene was purified from transformed *E. coli*, an animal cell expression vectors were constructed as TNFR/Fc fragment produced by restriction using EcoRI and XbaI was inserted at EcoRI/XbaI site of an animal cell expression vector, pCRTM3

(Invitrogen, USA) plasmid. And these were designated plasmid pTR11-Top10' and plasmid pTR22-Top10', and deposited as accession numbers of KCCM 10288 and KCCM 10291, respectively, at Korean Culture Center of Microorganisms (KCCM) on Jul. 10. 2001.

5 Transfection was performed by mixing either the plasmid pTR11-Top10' or plasmid pTR22-Top10' DNA including TNFR/Fc fusion genes as described above with the reagent of LipofectaminTM (Gibco BRL, USA). CHO-K1 cells with the concentration of $1\sim3 \times 10^5$ cells/well were inoculated in 6-well tissue culture plate (Nunc, USA), and incubated to 50~80% in 10% FBS - DMEM media, then the DNA-liposome complex, which was reacted for 15~45 min with 1~2 μ g of either the plasmid pTR11-Top10' or
10 plasmid pTR22-Top10' DNA including TNFR/Fc fusion genes as described above and 2~25 μ l of LipofectaminTM (Gibco BRL, USA), were added to the cell culture plate in the serum-free DMEM media. After incubation for 5 hrs, DMEM media with 20% serum was added and cells were incubated further for 18~24 hrs. After primary transfection, cells were incubated for 3 weeks in 10% FBS - DMEM media with 1.5mg/ml of Geneticin (G418,
15 Gibco BRL, USA), and formed colonies was selected for amplified incubation. The expression of fusion proteins was analyzed by ELISA using a peroxidase labeled goat anti-human IgG (KPL, USA).

ELISA was performed as follows. First, 1mg/ml of a peroxidase labeled goat anti-human IgG (KPL, USA) was diluted to 1:2,000 with 0.1M sodium bicarbonate, 100 μ l
20 of that was aliquoted into 96-well flexible plate (Falcon, USA) and sealed with plastic wrap, then incubated at 4 $^{\circ}$ C over 16 hrs to be coated on the surface of the plate. After this, it was washed for 3 times with washing buffer (0.1% Tween-20 in 1X PBS) and dilution buffer (48.5ml 1XPBS, 1.5ml FBS, 50 μ l Tween-20), and then was aliquoted to 180l. After 20 μ l of culture supernatant was dropped in the first well, then serially diluted using a
25 micropipette, and 0.01 μ g/ μ l of human immunoglobulin G (Sigma, USA) as the positive control and the culture media of untransfected CHO K-1 cell as the negative was equally diluted. After dilution, 96-well ELISA plate (Falcon, USA) was wrapped with aluminum

foil and incubated at 37°C for 1 hr 30 min, washed for 3 times with washing buffer. Peroxidase conjugated goat anti-human IgG (KPL, USA) was diluted to 1:5,000 with dilution buffer, aliquoted to 100µl, wrapped with aluminum foil, and reacted at 37°C for 1 hr. After reaction, this plate was washed for 3 times, colorized using TMB microwell
5 peroxidase substrate system (KPL, USA) and existence of expression was confirmed by measurement of absorbance at 655nm wavelength using microplate reader (Bio-Rad, Model 550, Japan).

Transfectants manufactured as above were designated TR11Ig-CHO and TR22Ig-CHO and deposited as accession numbers of KCLRF-BP-00046 and KCLRF-BP-
10 00049, respectively, at Korean Cell Line Research Foundation (KCLRF) on Jul. 7. 2001. And adaptation for transfectants as described above to one of the serum free media, CHO-S-SFM II (Gibco BRL, USA), was proceeded to purify the proteins produced by those transfectants as follows. After about 3×10^5 of cells were inoculated into the 6-well plate, cells were cultured at 5% CO₂, 37°C for over 16 hrs to adhere, and it was checked under a
15 microscope that cells were adhered at about 30~50% area of the plate, then cells were cultured in a media consisting of 10% FBS DMEM and CHO-S-SFM II in the ratio of 8:2. After culturing 3 times serial passage at this ratio, it was cultured 3 times at the ratio of 6:4; 3 times at 4:6; 3 times at 3:7; 3 times at 2:8; 3 times at 1:9; and finally cultured in 100% CHO-S-SFM II media. And the level of expression was measured by ELISA.

20 After these transfectant cells were cultured on a large scale in CHO-S-SFM II, the supernatants including each fusion proteins were centrifuged at 200X g for 12min to remove cell debris, and proteins were purified by the method using HiTrap protein A column (Amersham, USA) as follows. After 20mM of sodium phosphate (pH 7.0, Sigma, USA) was passed at the velocity of 1ml/min for 2 min, 10ml of supernatant was passed at
25 the same velocity to bind fusion protein to protein A. After 20mM of sodium phosphate (pH 7.0) was passed at the same velocity for 2 min to wash, 500µl of the extracts were serially fractionated in a 1.5ml tube as 0.1M of citric acid (pH 3.0, Sigma, USA) was

passed at the the same velocity for 3 min. This was adjusted to pH 7.0 using 1M of Tris (pH 11.0, USB, USA), the existence of fusion proteins in tube was confirmed through ELISA as described above. The purified proteins were concentrated by centrifugation at 2000Xg, 4 °C for 30min using Centricon 30 (Amicon, USA)

5

Example 5.

SDS-PAGE of purified TNFR1-TNFR1/Fc and TNFR2-TNFR2/Fc (Fig. 15)

Proteins purified using protein A column were electrophorized by the method of SDS-PAGE in reducing condition added by DTT, reducing reagent (which destroy disulfide bond), and in a non-reducing condition excluding DTT. The result of the estimation of molecular weight on SDS-PAGE is shown in Table 10. It was possible to confirm that TNFR/Fc proteins were the shape of a dimer in the cell. The molecular weight deduced from the amino acid sequence of TNFR1-TNFR1-Ig was about 70kDa, and was estimated as about 102kDa on SDS-PAGE. As this difference could be regarded as a general phenomenon which generate on the electrophoresis of glycoproteins, this feature seemed to occur as the result from decrease in mobility on the electrophoresis by the site of glycosylation.

15

20

Table 10. Molecular weight of TNFR-TNFR/Fc on the SDS-PAGE.

Proteins	Molecular weight (kDa)	
	Reducing condition	Non-reducing condition
TNFR1-TNFR1/Fc	102	200
TNFR2-TNFR2/Fc	115	220

Example 6.

Experiment of neutralization effect of simple/concatameric fusion dimeric TNFR/Fc fusion proteins on the cytotoxicity of TNF α and TNF β

5 An L929 cell [ATCC, *Mus musculus* (mouse), NCTC clone 929 (derivative of strain L; L-929; L cell) was used for testing the effect of TNFR/Fc fusion protein on the inhibition of cytotoxicity induced by TNF α and TNF β . This analysis was based on the TNFR activity of inhibiting cytotoxicity induced by TNF (Scallon et al., Cytokine 7:759, 1995).

10 L929 cells were inoculated to be 3×10^4 cells/well in 96-well plates, and incubated at 37°C for 24 hrs in a CO₂ incubator. Subsequently, actinomycin D (Sigma, USA) was added to 3 μ g/ml, and cells were incubated for 16~18 hrs with TNF α and TNF β in the concentration of expressing 100% cytotoxicity (0.5~2ng/ml), and with serially 10 times diluted TNFR sample. Then, the cells in the 96-well plate were stained by the staining reagent, crystal violet (Wako Pure Chemical Industries, Japan) and the activity of
15 the cells was estimated by the degree of absorbance at 595 nm wavelength using a spectrophotometer (Bio-Rad, Model-550, Japan).

As shown in Table 11 represented by IC₅₀ of each TNFR/Fc fusion protein, concatameric fusion proteins (TNFR1-TNFR1/Ig and TNFR2-TNFR2/Ig) have shown the higher inhibitory effect on the cytotoxicity induced by two kinds of TNF than simple
20 dimeric fusion proteins (TNFR1/Ig and TNFR2/Ig). Also, as compared with the effects of existing simple fusion dimer and concatameric shaped TNFR/Fc fusion protein dimer of the present invention on the inhibition of cytotoxicity of TNF α (Fig. 16) and TNF β (Fig. 17), it more clearly appeared that concatameric shaped TNFR/Fc fusion protein dimers of the present invention remarkably inhibited the TNF α and TNF β cytotoxicity.

25

Table 11. IC₅₀ of cytotoxicity inhibition

Fusion proteins	IC50 (ug/ml)
-----------------	--------------

		TNF α treated	TNF β treated
Simple dimer	[TNFR1/Fc] ₂	63	129
	[TNFR2/Fc] ₂	189	469
Concatameric dimer	[TNFR1-TNFR1/Fc] ₂	9	20
	[TNFR2-TNFR2/Fc] ₂	15	15

Example 7

Experiment of suppressive effect of simple/concatameric fusion dimeric CD2/Fc fusion protein and CTLA4/Fc fusion protein on the proliferation of active immune cell

5

WT100B1S, a cell line of B lymphocyte which was made by transfection of pyrexia patient's B lymphocyte with Epstein-Barr virus was incubated in RPMI 1640 supplemented with 10% FBS to use as antigen presenting cell of T lymphocyte. After centrifuged at 2,000rpm for 2 min to precipitate, this cells were resuspended in RPMI 1640 supplemented with 10% FBS to make 5.0×10^5 cells/ml, then irradiated by 3,000 rd of γ -ray.

10

T lymphocytes were isolated from blood of healthy adult using Ficoll-hypaque (Amersham, USA), then incubated RPMI 1640 supplemented with 10% FBS to 2.0×10^6 cells/ml.

15

To perform primary Mixed Lymphocyte Reaction (MLR), each 15ml of WT100B1S and T lymphocyte were mixed in 150mm cell culture dish, and incubated for 3 days, then added by 15ml of RPMI 1640 supplemented with 10% FBS and incubated for 3 days further. After incubated for total 6 days, live T lymphocytes were purified using Ficoll-hypaque (Amersham, USA) as described above, and purified T lymphocytes were stored in liquid nitrogen after freezing it by using the media comprising 45% FBS, 45% RPMI 1640, and 10% DMSO.

20

After T lymphocytes which were reacted by primary MLR were thawed to perform secondary MLR, the cells were washed with RPMI 1640 media for 2 times and made to be 3.0×10^5 cells/ml in RPMI 1640 supplemented with 10% FBS.

WT100B1S using as antigen presenting cell was newly cultured by the method as described above, then prepared by irradiation of 3,000 rd of γ -ray and to be 7.5×10^4 cells/ml in RPMI 1640 supplemented with 10% FBS. After 100 μ l of prepared WT100B1S was added in 96-well flat bottom cell culture plate and mixed with CD2/Fc and CTLA4/Fc fusion protein at final concentration of 10, 1, 10^{-1} , 10^{-2} , 10^{-3} , and 10^{-4} μ g/ml, 100 μ l of primary MLR reacted T lymphocytes as above was added. After incubated for 2 days in 5% CO₂, 37°C incubator, 100 μ l of RPMI 1640 supplemented with 10% FBS was added and incubated for 2 days further. In the last 6 hrs of the total 6 days culture, cells were incubated with addition of 1.2 μ Ci/ml of 3 H-thymidine (Amersham, USA).

At the end of culturing, supernatants were removed after centrifugation of 96-well plate was performed at 4°C, 110Xg for 10 min to precipitate T lymphocytes, and pellets were washed with 200 μ l of 1XPBS. Centrifugation was performed in the same condition and PBS was removed, then 200 μ l of ice-cold trichloridic acid (TCA, Merck, USA) was added and mixed for 2 min, then reacted at 4°C for 5 min to remove residue of 3 H-thymidine.

After centrifugation in the same condition as described above, supernatants were removed and T lymphocytes were fixed by incubation at 4°C for 5 min after 200 μ l of ice-cold 70% ethanol was added. Supernatants were removed after centrifugation, and 3 H-thymidine (Amersham, USA) residue was completely removed by treatment of 10% TCA in the same method as described above.

Cell lysis was performed by reaction with 100 μ l of 2% SDS (pH 8.0) and 0.5N of NaOH at 37°C for 30min, and T lymphocytes were precipitated by centrifugation at 25°C, 110Xg for 10min, and then 50 μ l of supernatants was transferred to 96-well sample plate (Wallac, USA). After 1.5 volume of OptiPhase SuperMix (Wallac, USA) was added into the supernatants and mixed for 5 min, the existence of T lymphocyte proliferation was confirmed by measurement of cpm value of 3 H using 1450 MicroBeta TriLux microplate liquid scintillation andluminescence counter (Wallac, USA).

Example 8

Experiment of effect on increase of plasma half-life of glycosylated concatameric fusion dimeric proteins in mouse

5 The measurement of plasma half-life of glycosylated concatameric fusion dimeric proteins, [mgTNFR1-TNFR1/Fc]2, [mgTNFR2-TNFR2/Fc]2, [mgCD2-CD2/Fc]2, and [mgCTLA4-CTLA4/Fc]2 was performed by measuring the concentration of proteins using ELISA after 5µg of purified fusion proteins was i.p. injected into mouse (ICR, 10 Samtako, Korea) and bloods were extracted at regular interval for 120 hrs (5 days) as maximum. As shown Fig. 20, Fig. 21, and Fig 22, it could be seen that the plasma half-life of glycosylated concatameric fusion dimeric proteins have been increased in comparison of the corresponding simple fusion dimeric proteins of native shape, and the increase in efficacy through continuous effect could be expected.

Example 9

Experiment of effects of simple/concatameric TNFR/Fc fusion protein dimers on collagen-induced arthritis of DBA/1 mouse

20 Collagen Induced Arthritis (CIA) was developed by injection with 100µg per DBA/1 mouse of type II collagen dissolved at 2mg/ml concentration in 0.05M acetic acid and Arthrogen-CIA adjuvant (Chondrex, USA) into tail. Boosting was performed after 3 weeks, and incomplete Freund's adjuvant (Difco, USA) was used.

25 Arthritis was developed 3~4 weeks after immunization with 100µg of type II collagen in the DBA/1 mice. Red and swollen paws of mice had been observed 3~5 days after onset, and inflammatory arthritis lasted more than 3 - 4 weeks. Although inflammation was eventually alleviated, damaged joints remained rigid permanently. The degree of

arthritis was measured 2~3 times per week on the basis of table 12 which represented subjective index of arthritis severity (measure average of five mice in each experiment). To measure the effects of simple and concatameric fusion dimeric TNFR/Fc on CIA, TNFR/Fc or PBS was i.p. injected into the mice. TNFR/Fc was injected with 10 μ g at every 2 days for 19~45 days into 5 mice per experiments (arrows in Fig. 23). PBS was injected into 5 mice as control. As shown in Fig. 7, in the case of mice injected with existing simple dimeric shaped TNFR/Fc fusion protein, it could be seen that the effect decreased to about 26-38% in comparison with the figures of arthritis index in mice injected with PBS as control, but 42-55% decreased in case of concatameric shaped dimer, [TNFR1-TNFR1/Fc]₂ and [TNFR2-TNFR2/Fc]₂ were injected. Therefore, it could be shown that concatameric fusion dimeric TNFR/Fc fusion proteins have remarkably decreased arthritis of mouse than existing simple fusion dimeric TNFR/Fc fusion proteins.

Table 12. Severity score of arthritis

Severity score	Condition of disease
0	No erythema and swelling
1	Erythema and mild swelling limited to ankle and tarsal
2	Erythema and mild swelling spread from ankle to tarsal
3	Erythema and mild swelling spread from ankle to metatarsal joint
4	Erythema and severe swelling extend to ankle, legs, and digits

The results as above represented that concatameric shaped dimeric TNFR/Fc fusion proteins were more effective in decreasing the rate of CIA development than existing simple dimeric fusion proteins, therefore, as use in arthritis therapy, concatameric shaped protein compositions could be more effective therapeutics than existing protein compositions.

The concatameric proteins, concatameric fusion dimeric proteins and their glycosylated proteins of the present invention were able to express increased efficacy and high stability, and to be produced with high yield.

5

INDUSTRIAL APPLICABILITY

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 28, line 0-5	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korea Cell Line Research Foundation(KCLRF)	
Address of depositary institution(including postal code and country) Cancer Research Institute, Seoul National University College of Medicine 28 Yongon-dong, Chongno-gu SEOUL 120-091 Republic of Korea	
Date of deposit 29/06/2001	Accession Number KCLRF-BP-00046
C.ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D.DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E.SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

Form PCT/RO/134(July 1998)

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 28, line 0-5	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korea Cell Line Research Foundation(KCLRF)	
Address of depositary institution(including postal code and country) Cancer Research Institute, Seoul National University College of Medicine 28 Yongon-dong, Chongno-gu SEOUL 120-091 Republic of Korea	
Date of deposit 29/06/2001	Accession Number KCLRF-BP-00049
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line 15-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 22/07/2002	Accession Number KCCM 10407
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line 15-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10399
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For international Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line 51-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10401
C.ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D.DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E.SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 27, line 10-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10400
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For international Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 27, line 10-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10402
C.ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D.DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E.SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line <u>15-20</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution <i>(including postal code and country)</i> 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10404
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line 15-20 .	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10403
C.ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D.DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E.SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For international Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 29, line 15-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are on an additional sheet <input type="checkbox"/>	
Name of depositary institution Korean Culture Center of Microorganisms(KCCM)	
Address of depositary institution(including postal code and country) 361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, SEOUL 120-091, Republic of Korea	
Date of deposit 11/07/2002	Accession Number KCCM 10405
C.ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D.DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E.SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For international Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer
---	--

WHAT IS CLAIMED IS:

1. A concatameric protein comprising two soluble domains, in which a N-terminus of a soluble domain of a biologically active protein is linked to C-terminus of an identical soluble domain or a different soluble domain of a biologically active protein.
- 5 2. A concatameric fusion dimeric protein comprising two monomeric proteins formed by linkage of a concatamer of two identical soluble extracellular domains of proteins involving immune response to a hinge region of an Fc fragment of an immunoglobulin molecule, wherein said monomeric proteins are linked by intermolecular disulfide bonds at the hinge region, and having improved stability and therapeutic effects.
- 10 3. The concatameric fusion dimeric protein as set forth in claim 2, wherein the immunoglobulin molecule is IgG.
4. The concatameric fusion dimeric protein as set forth in claim 2, wherein the protein involving immune response is selected from the group consisting of cytokines, cytokine receptors, adhesion molecules, tumor necrosis factor receptors, receptor tyrosine
15 kinases, chemokine receptors and other cell surface proteins which contain a soluble extracellular domain.
5. The concatameric fusion dimeric protein as set forth in claim 4, wherein the protein is selected from the group consisting of IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12, IL-17, TNF, TGF, IFN, GM-CSF, G-CSF, EPO, TPO, M-CSF, GHR, IL-13R, IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-9R, IL-15R, TNFR, TGFR, IFNR, interferon- α R, - β R and - γ R, GM-CSFR, G-CSFR, EPOR, cMpl, gp130, Fas (Apo 1), CCR1, CXCR1-4, TrkA, TrkB, TrkC, Htk, REK7, Rse/Tyro-3, hepatocyte growth factor R, platelet-derived growth factor R, Flt-1, CD2, CD4, CD5, CD6, CD22, CD27, CD28, CD30, CD31, CD40, CD44, CD100, CD137, CD150, LAG-3, B7, B61, β -neurexin, CTLA-4,
20 IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-9R, IL-15R, TNFR, TGFR, IFNR, interferon- α R, - β R and - γ R, GM-CSFR, G-CSFR, EPOR, cMpl, gp130, Fas (Apo 1), CCR1, CXCR1-4, TrkA, TrkB, TrkC, Htk, REK7, Rse/Tyro-3, hepatocyte growth factor R, platelet-derived growth factor R, Flt-1, CD2, CD4, CD5, CD6, CD22, CD27, CD28, CD30, CD31, CD40, CD44, CD100, CD137, CD150, LAG-3, B7, B61, β -neurexin, CTLA-4,
25 ICOS, ICAM-1, complement R-2 (CD21), IgER, lysosomal membrane gp-1, α 2-microglobulin receptor-related proteins, and sodium-releasing peptide R.

6. The concatameric fusion dimeric protein as set forth in claim 2, wherein the monomeric protein contains an amino acid sequence of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 18, or SEQ ID NO: 20.

7. A DNA construct encoding a monomeric protein formed by linkage of a concatamer of two identical soluble extracellular domains of a protein involving immune response to a hinge region of an Fc fragment of an immunoglobulin molecule.

8. The DNA construct as set forth in claim 7, wherein the immunoglobulin molecule is IgG.

9. The DNA construct as set forth in claim 7, wherein the protein involving immune response is selected from the group consisting of cytokines, cytokine receptors, adhesion molecules, tumor necrosis factor receptors, receptor tyrosine kinases, chemokine receptors and other cell surface proteins which contain a soluble extracellular domain.

10. The DNA construct as set forth in claim 9, wherein the protein is selected from the group consisting of IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12, IL-17, TNF, TGF, IFN, GM-CSF, G-CSF, EPO, TPO, M-CSF, GHR, IL-13R, IL-1R, IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-9R, IL-15R, TNFR, TGFR, IFNR, interferon- α R, - β R and - γ R, GM-CSFR, G-CSFR, EPOR, cMpl, gp130, Fas (Apo 1), CCR1, CXCR1-4, TrkA, TrkB, TrkC, Htk, REK7, Rse/Tyro-3, hepatocyte growth factor R, platelet-derived growth factor R, Flt-1, CD2, CD4, CD5, CD6, CD22, CD27, CD28, CD30, CD31, CD40, CD44, CD100, CD137, CD150, LAG-3, B7, B61, β -neurexin, CTLA-4, ICOS, ICAM-1, complement R-2 (CD21), IgER, lysosomal membrane gp-1, α 2-microglobulin receptor-related proteins, and sodium-releasing peptide R.

11. The DNA construct as set forth in claim 7, wherein the DNA construct contains a nucleotide sequence of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 17, or SEQ ID NO: 19.

12. A recombinant expression plasmid comprising the DNA construct of claim 7 operably linked thereto.

13. The recombinant expression plasmid as set forth in claim 12, wherein the recombinant expression plasmid is a pTR11-Top10' plasmid (accession No.: KCCM 10288),
5 a pTR22-Top10' plasmid (accession No.: KCCM 10289), a pCD22Ig plasmid (accession No.: KCCM 10402), or a pCT44Ig plasmid (accession No.: KCCM 10400).

14. A host cell transformed or transfected with the recombinant expression plasmid of claim 12.

15. The host cell as set forth in claim 14, wherein the host cell is a
10 mammalian cell.

16. The host cell as set forth in claim 14 or 15, wherein the recombinant expression plasmid is a pTR11-Top10' plasmid (accession No.: KCCM 10288), a pTR22-Top10' plasmid (accession No.: KCCM 10289), a pCD22Ig plasmid (accession No.: KCCM 10402), or a pCT44Ig plasmid (accession No.: KCCM 10400).

17. The host cell as set forth in claim 16, wherein the host cell is a TR11Ig-CHO cell line (accession No.: KCLRF-BP-00046) or a TR22Ig-CHO cell line (accession No.: KCLRF-BP-00049).

18. A method of preparing a concatameric fusion dimeric protein in which disulfide bonds are formed between the hinge regions of two monomeric proteins,
20 comprising the steps of:

culturing the transformed or transfected host cell of claim 14 under conditions suitable for expression of a DNA construct encoding a concatameric fusion monomeric protein in which a concatamer of two identical soluble extracellular domains of

proteins involving immune response is linked to a hinge region of an Fc fragment of an immunoglobulin molecule; and

isolating and purifying a dimeric protein formed by dimerization of the produced monomeric proteins from culture medium.

5 19. The method as set forth in claim 18, wherein the DNA construct encoding a concatameric fusion monomeric protein is prepared by preparing a DNA construct encoding a simple fusion monomeric protein formed by joining a DNA fragment encoding an Fc fragment of an immunoglobulin molecule and a DNA fragment encoding a soluble extracellular domain of a protein involving immune response; and joining the prepared
10 DNA construct and a second DNA fragment identical to the DNA fragment encoding a soluble extracellular domain of a protein involving immune response.

20. The method as set forth in claim 19, wherein the DNA construct encoding a concatameric fusion monomeric protein contains a glycosylation motif sequence.

15 21. The method as set forth in claim 20, wherein the glycosylation motif sequence is inserted to a region at which two soluble extracellular domains are joined.

22. The method as set forth in claim 19, wherein the concatameric fusion monomeric protein contains a leader sequence.

20 23. The method as set forth in claim 22, wherein the concatameric fusion monomeric protein is CTLA-4, and the leader sequence has an amino acid sequence of MACLGFRHKAQKNLAARTWPCTLLFFIPVFCKA.

24. The method as set forth in claim 23, wherein the leader sequence has an amino acid sequence of MRTWPCTLLFFIPVFCKA excluding ACLGFRHKAQKNLAA.

25. The method as set forth in any of claims 18 to 24, wherein the host cell is a mammalian cell.

26. A concatameric fusion dimeric protein comprising two monomeric proteins formed by linkage of a concatamer of two identical soluble extracellular domains of proteins involving immune response to the hinge region of Fc fragment of an immunoglobulin molecule, wherein said monomeric proteins are linked by formation of intermolecular disulfide bonds at the hinge region and glycosylated, and having improved stability and therapeutic effects.

27. The concatameric fusion dimeric protein as set forth in claim 26, wherein the monomeric protein contains an amino acid sequence of SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 22, or SEQ ID NO: 24.

28. A DNA construct encoding a monomeric protein formed by linkage of a concatamer of two identical soluble extracellular domains of proteins involving immune response to a hinge region of an Fc fragment of an immunoglobulin molecule and containing glycosylation motif peptides.

29. The DNA construct as set forth in claim 28, wherein the DNA construct contains an amino acid sequence of SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 21, or SEQ ID NO: 23.

30. A recombinant expression plasmid operably linked to the DNA construct of claim 28.

31. The recombinant expression plasmid as set forth in claim 30, wherein the recombinant expression plasmid is a pTR11Ig-MG plasmid (accession No.: KCCM 10404), a pTR22Ig-MG plasmid (accession No.: KCCM 10407), a pCD22Ig-MG plasmid (accession No.: KCCM 10401), or a pCT44Ig-MG plasmid (accession No.: KCCM 10399).

32. A host cell transformed or transfected with the recombinant expression plasmid of claim 30.


33. The host cell as set forth in claim 32, wherein the host cell is a mammalian cell.

34. A pharmaceutical or diagnostic composition comprising the dimeric protein of claim 2.

5 35. A pharmaceutical or diagnostic composition comprising the glycosylated dimeric protein of claim 26.


1/23

FIG. 1



2/23


FIG. 2




3/23

FIG. 3

a




b




4/23

FIG. 4




5/23

FIG. 5




6/23

FIG. 6




7/23

FIG. 7




8/23

FIG. 8




9/23

FIG. 9




10/23

FIG. 10




11/23

FIG. 11




12/23

FIG. 12




13/23

FIG. 13




14/23

FIG. 14




15/23

FIG. 15




16/23

FIG. 16




17/23

FIG. 17




18/23

FIG. 18




19/23

FIG. 19




20/23

FIG. 20




21/23

FIG. 21




22/23

FIG. 22



23/23

FIG. 23



Sequence Listing

- <110> MeDexGen Inc.
CHUNG, Yong Hoon
HAN, Ji Woong
LEE, Hye Ja
CHOI, Eun Yong
KIM, Jin Mi
YIM, Soo Bin
- <120> Method of manufacturing Ig-fusion proteins by concatamerization,
TNFR/Fc, CD2/Fc, CTLA4/Fc fusion proteins manufactured by the
method, DNA coding the proteins, vectors including the DNA, and
cells transformed by the vectorTOR
- <160> 52
- <170> KopatentIn 1.71
- <210> 1
<211> 1335
<212> DNA
<213> Homo sapiens
- <220>
<221> CDS
<222> (1)..(1332)
<223> TNFR1-IgG
- <220>
<221> C_region
<222> (634)..(1335)
<223> Hinge, CH2, CH3 region
- <220>
<221> misc_signal
<222> (160)..(168)
<223> N-linked glycosylation site

Sequence Listing

<220>

<221> misc_signal

<222> (433)..(441)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (451)..(459)

<223> N-linked glycosylation site

<220>

<221> primer_bind

<222> (1)..(15)

<223> PCR primer SEQ ID : 25 binding site

<220>

<221> primer_bind

<222> (616)..(652)

<223> PCR primer SEQ ID : 26(antisense) binding site

<220>

<221> primer_bind

<222> (616)..(651)

<223> PCR primer SEQ ID : 27 binding site

<220>

<221> primer_bind

<222> (1312)..(1335)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(60)

<223> signal peptide

Sequence Listing

<400> 1
 atg ggc ctc tcc acc gtg cct gac ctg ctg ctg ccg ctg gtg ctc ctg 48
 Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu
 1 5 10 15

gag ctg ttg gtg gga ata tac ccc tca ggg gtt att gga ctg gtc cct 96
 Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro
 20 25 30

cac cta ggg gac agg gag aag aga gat agt gtg tgt ccc caa gga aaa 144
 His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys
 35 40 45

tat atc cac cct caa aat aat tcg att tgc tgt acc aag tgc cac aaa 192
 Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys
 50 55 60

gga acc tac ttg tac aat gac tgt cca ggc ccg ggg cag gat acg gac 240
 Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp
 65 70 75 80

tgc agg gag tgt gag agc ggc tcc ttc acc gct tca gaa aac cac ctc 288
 Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu
 85 90 95

aga cac tgc ctc agc tgc tcc aaa tgc cga aag gaa atg ggt cag gtg 336
 Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val
 100 105 110

gag atc tct tct tgc aca gtg gac cgg gac acc gtg tgt ggc tgc agg 384
 Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg
 115 120 125

aag aac cag tac cgg cat tat tgg agt gaa aac ctt ttc cag tgc ttc 432
 Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe
 130 135 140

aat tgc agc ctc tgc ctc aat ggg acc gtg cac ctc tcc tgc cag gag 480

Asn Cys Ser Leu Cys	Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu		
145	150	155	160
aaa cag aac acc gtg tgc acc tgc cat gca ggt ttc ttt cta aga gaa		528	
Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu			
165	170	175	
aac gag tgt gtc tcc tgt agt aac tgt aag aaa agc ctg gag tgc acg		576	
Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr			
180	185	190	
aag ttg tgc cta ccc cag att gag aat gtt aag ggc act gag gac tca		624	
Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser			
195	200	205	
ggc acc aca gca gag ccc aaa tct tgt gac aaa act cac aca tgc cca		672	
Gly Thr Thr Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro			
210	215	220	
cgg tgc cca gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc		720	
Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe			
225	230	235	240
ccc cca aaa ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc		768	
Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val			
245	250	255	
aca tgc gtg gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc		816	
Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe			
260	265	270	
aac tgg tac gtg gac gcc gtg gag gtg cat aat gcc aag aca aag ccg		864	
Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro			
275	280	285	
cgg gag gag cag tac aac agc acg tac cgg gtg gtc agc gtc ctc acc		912	
Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr			
290	295	300	
gtc ctg cac caq gac tqg ctg aat gcc aag gag tac aag tgc aaq gtc		960	

Sequence Listing

Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val		
305						310				315					320		
tcc	aac	aaa	gcc	ctc	cca	gcc	ccc	atc	gag	aaa	acc	atc	toc	aaa	gcc		1008
Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala		
			325					330					335				
aaa	ggg	cag	ccc	cga	gaa	cca	cag	gtg	tac	acc	ctg	ccc	cca	tcc	cgg		1056
Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg		
			340					345					350				
gat	gag	ctg	acc	aag	aac	cag	gtc	agc	ctg	acc	tgc	ctg	gtc	aaa	ggc		1104
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly		
		355					360					365					
ttc	tat	ccc	agc	gac	atc	gcc	gtg	gag	tgg	gag	agc	aat	ggg	cag	ccg		1152
Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro		
	370					375				380							
gag	aac	aac	tac	aag	acc	acg	cct	ccc	gtg	ctg	gac	tcc	gac	ggc	tcc		1200
Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser		
385				390					395				400				
tcc	ttc	ctc	tac	agc	aag	ctc	acc	gtg	gac	aag	agc	agg	tgg	cag	cag		1248
Ser	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln		
			405					410					415				
ggg	aac	gtc	ttc	tca	tgc	tcc	gtg	atg	cat	gag	gct	ctg	cac	aac	cac		1296
Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His		
		420						425					430				
tac	acg	cag	aag	agc	ctc	tcc	ctg	tct	ccg	ggt	aaa		tga				1335
Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys						
		435						440									

<210> 2

<211> 444

<212> PRT

<213> Homo sapiens

Sequence Listing

<400> 2

Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu

1 5 10 15

Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro

20 25 30

His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys

35 40 45

Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys

50 55 60

Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp

65 70 75 80

Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu

85 90 95

Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val

100 105 110

Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg

115 120 125

Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe

130 135 140

Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu

145 150 155 160

Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu

165 170 175

Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr

180 185 190

Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser

195 200 205

Sequence Listing

Gly Thr Thr Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro
 210 215 220
 Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe
 225 230 235 240
 Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 245 250 255
 Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe
 260 265 270
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 275 280 285
 Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
 290 295 300
 Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 305 310 315 320
 Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala
 325 330 335
 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 340 345 350
 Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 355 360 365
 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 370 375 380
 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
 385 390 395 400
 Ser Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
 405 410 415

Sequence Listing

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 420 425 430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440

<210> 3
 <211> 1473
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1470)
 <223> TNFR2-IgG

<220>
 <221> C_region
 <222> (772)..(1473)
 <223> Hinge, CH2, CH3 region

<220>
 <221> misc_signal
 <222> (511)..(519)
 <223> N-linked glycosylation site

<220>
 <221> misc_signal
 <222> (577)..(585)
 <223> N-linked glycosylation site

<220>
 <221> primer_bind
 <222> (1)..(15)
 <223> PCR primer SEQ ID : 29 binding site

Sequence Listing

<220>
 <221> primer_bind
 <222> (754)..(790)
 <223> PCR primer SEQ ID : 30(antisense) binding site

<220>
 <221> primer_bind
 <222> (754)..(790)
 <223> PCR primer SEQ ID : 31 binding site

<220>
 <221> primer_bind
 <222> (1451)..(1473)
 <223> PCR primer SEQ ID : 28(antisense) binding site

<220>
 <221> sig_peptide
 <222> (1)..(66)
 <223> signal peptide

<400>	3	
atg gcg ccc gtc gcc gtc tgg gcc gcg ctg gcc gtc gga ctg gag ctc		48
Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu		
1 5 10 15		
tgg gct gcg gcg cac gcc ttg ccc gcc cag gtg gca ttt aca ccc tac		96
Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr		
20 25 30		
gcc ccg gag ccc ggg agc aca tgc cgg ctc aga gaa tac tat gac cag		144
Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln		
35 40 45		
aca gct cag atg tgc tgc agc aaa tgc tgc ccg gcc caa cat gca aaa		192

Sequence Listing

Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys	
50 55 60	
gtc ttc tgt acc aag acc tcg gac acc gtg tgt gac tcc tgt gag gac	240
Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp	
65 70 75 80	
agc aca tac acc cag ctc tgg aac tgg gtt ccc gag tgc ttg agc tgt	288
Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys	
85 90 95	
ggc tcc cgc tgt agc tct gac cag gtg gaa act caa gcc tgc act cgg	336
Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg	
100 105 110	
gaa cag aac cgc atc tgc acc tgc agg ccc ggc tgg tac tgc gcg ctg	384
Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu	
115 120 125	
agc aag cag gag ggg tgc cgg ctg tgc gcg ccg ctg cgc aag tgc cgc	432
Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg	
130 135 140	
ccg ggc ttc ggc gtg gcc aga cca gga act gaa aca tca gac gtg gtg	480
Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val	
145 150 155 160	
tgc aag ccc tgt gcc ccg ggg acg ttc tcc aac acg act tca tcc acg	528
Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr	
165 170 175	
gat att tgc agg ccc cac cag atc tgt aac gtg gtg gcc atc cct ggg	576
Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly	
180 185 190	
aat gca agc atg gat gca gtc tgc acg tcc acg tcc ccc acc cgg agt	624
Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser	
195 200 205	
atg gcc cca ggg gca gta cac tta ccc cag cca gtg tcc aca cga tcc	672

Sequence Listing

Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser	
210	215 220
caa cac acg cag cca act cca gaa ccc agc act gct cca agc acc tcc	720
Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser	
225	230 235 240
ttc ctg ctc cca atg ggc ccc agc ccc cca gct gaa ggg agc act ggc	768
Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly	
245	250 255
gac gca gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg tgc	816
Asp Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys	
260	265 270
cca gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca	864
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro	
275	280 285
aaa ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc aca tgc	912
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys	
290	295 300
gtg gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg	960
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp	
305	310 315 320
tac gtg gac ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg gag	1008
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu	
325	330 335
gag cag tac aac agc acg tac cgg gtg gtc agc gtc ctc acc gtc ctg	1056
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu	
340	345 350
cac cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc aac	1104
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn	
355	360 365
aaa gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg	1152

Sequence Listing

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 370 375 380

cag ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat gag 1200
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
 385 390 395 400

ctg acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat 1248
 Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 405 410 415

ccc agc gac atc gcc gtg gag tgg gag agc aat ggg cag ccg gag aac 1296
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 420 425 430

aac tac aag acc acg cct ccc gtg ctg gac tcc gac ggc tcc tcc ttc 1344
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Ser Phe
 435 440 445

ctc tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac 1392
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 450 455 460

gtc ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg 1440
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 465 470 475 480

cag aag agc ctc tcc ctg tct ccg ggt aaa tga 1473
 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 485 490

<210> 4

<211> 490

<212> PRT

<213> Homo sapiens

<400> 4

Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15

Sequence Listing

Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45
 Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60
 Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
 65 70 75 80
 Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
 85 90 95
 Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
 100 105 110
 Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
 115 120 125
 Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
 130 135 140
 Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
 145 150 155 160
 Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
 165 170 175
 Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
 180 185 190
 Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
 195 200 205
 Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
 210 215 220

Sequence Listing

```

Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
225                      230                      235                      240

Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly
                      245                      250                      255

Asp Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
                      260                      265                      270

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
                      275                      280                      285

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
                      290                      295                      300

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
305                      310                      315                      320

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
                      325                      330                      335

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
                      340                      345                      350

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
                      355                      360                      365

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
                      370                      375                      380

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
385                      390                      395                      400

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
                      405                      410                      415

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
                      420                      425                      430

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Ser Phe

```

Sequence Listing

```

      435      440      445
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      450      455      460
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      465      470      475      480
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      485      490

```

```

<210> 5
<211> 1887
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (1)..(1884)
<223> TNFR1-TNFR1-IgG

```

```

<220>
<221> C_region
<222> (1716)..(1887)
<223> Hinge, CH2, CH3 region

```

```

<220>
<221> misc_signal
<222> (160)..(168)
<223> N-linked glycosylation site

```

```

<220>
<221> misc_signal
<222> (433)..(441)
<223> N-linked glycosylation site

```


Sequence Listing

<220>
<221> misc_signal
<222> (451)..(459)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (631)..(639)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (712)..(720)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (985)..(993)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (1003)..(1011)
<223> N-linked glycosylation site

<220>
<221> primer_bind
<222> (1)..(15)
<223> PCR primer SEQ ID : 25 binding site

<220>
<221> primer_bind
<222> (592)..(628)

Sequence Listing

<223> PCR primer SEQ ID : 33(antisense) binding site

<220>

<221> primer_bind

<222> (622)..(655)

<223> PCR primer SEQ ID : 32 binding site

<220>

<221> primer_bind

<222> (1168)..(1204)

<223> PCR primer SEQ ID : 26(antisense) binding site

<220>

<221> primer_bind

<222> (1168)..(1204)

<223> PCR primer SEQ ID : 27 binding site

<220>

<221> primer_bind

<222> (1864)..(1887)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(60)

<223> signal peptide

<400> 5

atg ggc ctc tcc acc gtg cct gac ctg ctg ctg ccg ctg gtg ctc ctg 48
Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu
1 5 10 15

gag ctg ttg gtg gga ata tac ccc tca ggg gtt att gga ctg gtc cct 96
Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro

Sequence Listing

20	25	30	
cac cta ggg gac agg gag aag aga gat agt gtg tgt ccc caa gga aaa			144
His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys			
35	40	45	
tat atc cac cct caa aat aat tcg att tgc tgt acc aag tgc cac aaa			192
Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys			
50	55	60	
gga acc tac ttg tac aat gac tgt cca ggc cgg ggg cag gat acg gac			240
Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp			
65	70	75	80
tgc agg gag tgt gag agc ggc tcc ttc acc gct tca gaa aac cac ctc			288
Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu			
85	90	95	
aga cac tgc ctc agc tgc tcc aaa tgc cga aag gaa atg ggt cag gtg			336
Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val			
100	105	110	
gag atc tct tct tgc aca gty gac cgg gac acc gtg tgt ggc tgc agg			384
Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg			
115	120	125	
aag aac cag tac cgg cat tat tgg agt gaa aac ctt ttc cag tgc ttc			432
Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe			
130	135	140	
aat tgc agc ctc tgc ctc aat ggg acc gtg cac ctc tcc tgc cag gag			480
Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu			
145	150	155	160
aaa cag aac acc gtg tgc acc tgc cat gca ggt ttc ttt cta aga gaa			528
Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu			
165	170	175	
aac gag tgt gtc tcc tgt agt aac tgt aag aaa agc ctg gag tgc acg			576
Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr			

Sequence Listing

180	185	190	
aag ttg tgc cta ccc cag att gag aat gtt aag ggc act gag gac gga			624
Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Gly			
195	200	205	
tcc ggg aac att tca ctg gtc cct cac cta ggg gac agg gag aag aga			672
Ser Gly Asn Ile Ser Leu Val Pro His Leu Gly Asp Arg Glu Lys Arg			
210	215	220	
gat agt gtg tgt ccc caa gga aaa tat atc cac cct caa aat aat tcg			720
Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser			
225	230	235	240
att tgc tgt acc aag tgc cac aaa gga acc tac ttg tac aat gac tgt			768
Ile Cys Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys			
245	250	255	
cca ggc ccg ggg cag gat acg gac tgc agg gag tgt gag agc ggc tcc			816
Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser			
260	265	270	
ttc acc gct tca gaa aac cac ctc aga cac tgc ctc agc tgc tcc aaa			864
Phe Thr Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys			
275	280	285	
tgc cga aag gaa atg ggt cag gtg gag atc tct tct tgc aca gtg gac			912
Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp			
290	295	300	
cgg gac acc gtg tgt ggc tgc agg aag aac cag tac cgg cat tat tgg			960
Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp			
305	310	315	320
agt gaa aac ctt ttc cag tgc ttc aat tgc agc ctc tgc ctc aat ggg			1008
Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly			
325	330	335	
acc gtg cac ctc tcc tgc cag gag aaa cag aac acc gtg tgc acc tgc			1056
Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys			

Sequence Listing

340	345	350	
cat gca ggt ttc ttt cta aga gaa aac gag tgt gtc tcc tgt agt aac			1104
His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser Asn			
355	360	365	
tgt aag aaa agc ctg gag tgc acg aag ttg tgc cta ccc cag att gag			1152
Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu			
370	375	380	
aat gtt aag ggc act gag gac tca ggc acc aca gca gag ccc aaa tot			1200
Asn Val Lys Gly Thr Glu Asp Ser Gly Thr Thr Ala Glu Pro Lys Ser			
385	390	395	400
tgt gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa ctc ctg			1248
Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu			
405	410	415	
ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc			1296
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu			
420	425	430	
atg atc tcc ccg acc cct gag gtc aca tgc gtg gtg gtg gac gtg agc			1344
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser			
435	440	445	
cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag			1392
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu			
450	455	460	
gtg cat aat gcc aag aca aag ccg ccg gag gag cag tac aac agc acg			1440
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr			
465	470	475	480
tac ccg gtg gtc agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat			1488
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn			
485	490	495	
ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc			1536
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro			

Sequence Listing

500	505	510	
atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa cca cag			1584
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln			
515	520	525	
gtg tac acc ctg ccc cca tcc cgg gat gag ctg acc aag aac cag gtc			1632
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val			
530	535	540	
agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc gcc gtg			1680
Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val			
545	550	555	560
gag tgg gag agc aat ggg cag ccg gag aac aac tac aag acc acg cct			1728
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro			
565	570	575	
ccc gtg ctg gac tcc gac ggc tcc tcc ttc ctc tac agc aag ctc acc			1776
Pro Val Leu Asp Ser Asp Gly Ser Ser Phe Leu Tyr Ser Lys Leu Thr			
580	585	590	
gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc tcc gtg			1824
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val			
595	600	605	
atg cat gag gct ctg cac aac cac tac acg cag aag agc ctc tcc ctg			1872
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu			
610	615	620	
tct ccg ggt aaa	tga		1887
Ser Pro Gly Lys			
625			
<210> 6			
<211> 628			
<212> PRT			
<213> Homo sapiens			

Sequence Listing

<400> 6

Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu
 1 5 10 15

Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro
 20 25 30

His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys
 35 40 45

Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys
 50 55 60

Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp
 65 70 75 80

Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu
 85 90 95

Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val
 100 105 110

Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg
 115 120 125

Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe
 130 135 140

Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu
 145 150 155 160

Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu
 165 170 175

Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr
 180 185 190

Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Gly
 195 200 205

Sequence Listing

Ser Gly Asn Ile Ser Leu Val Pro His Leu Gly Asp Arg Glu Lys Arg
 210 215 220

Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser
 225 230 235 240

Ile Cys Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys
 245 250 255

Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser
 260 265 270

Phe Thr Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys
 275 280 285

Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp
 290 295 300

Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp
 305 310 315 320

Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly
 325 330 335

Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys
 340 345 350

His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser Asn
 355 360 365

Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu
 370 375 380

Asn Val Lys Gly Thr Glu Asp Ser Gly Thr Thr Ala Glu Pro Lys Ser
 385 390 395 400

Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
 405 410 415

Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu

Sequence Listing

420	425	430
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser		
435	440	445
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu		
450	455	460
Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr		
465	470	475
		480
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn		
485	490	495
Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro		
500	505	510
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln		
515	520	525
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val		
530	535	540
Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val		
545	550	555
		560
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro		
565	570	575
Pro Val Leu Asp Ser Asp Gly Ser Ser Phe Leu Tyr Ser Lys Leu Thr		
580	585	590
Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val		
595	600	605
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu		
610	615	620
Ser Pro Gly Lys		
625		

Sequence Listing

<210> 7
<211> 2163
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(2160)
<223> TNFR2-TNFR2-IgG

<220>
<221> C_region
<222> (1462)..(2163)
<223> Hinge, CH2, CH3 region

<220>
<221> misc_signal
<222> (511)..(519)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (577)..(585)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (769)..(777)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (1201)..(1209)

Sequence Listing

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (1267)..(1275)

<223> N-linked glycosylation site

<220>

<221> primer_bind

<222> (1)..(15)

<223> PCR primer SEQ ID : 29 binding site

<220>

<221> primer_bind

<222> (761)..(795)

<223> PCR primer SEQ ID : 35(antisense) binding site

<220>

<221> primer_bind

<222> (741)..(768)

<223> PCR primer SEQ ID : 34 binding site

<220>

<221> primer_bind

<222> (1444)..(1480)

<223> PCR primer SEQ ID : 30(antisense) binding site

<220>

<221> primer_bind

<222> (1444)..(1480)

<223> PCR primer SEQ ID : 31 binding site

<220>

Sequence Listing

<221> primer_bind
 <222> (2141)..(2163)
 <223> PCR primer SEQ ID : 28(antisense) binding site

<220>
 <221> sig_peptide
 <222> (1)..(66)
 <223> signal peptide

<400> 7
 atg gcg ccc gtc gcc gtc tgg gcc gcg ctg gcc gtc gga ctg gag ctc 48
 Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15
 tgg gct gcg gcg cac gcc ttg ccc gcc cag gtg gca ttt aca ccc tac 96
 Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 gcc ccg gag ccc ggg agc aca tgc cgg ctc aga gaa tac tat gac cag 144
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45
 aca gct cag atg tgc tgc agc aaa tgc tcg ccg ggc caa cat gca aaa 192
 Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60
 gtc ttc tgt acc aag acc tcg gac acc gtg tgt gac tcc tgt gag gac 240
 Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
 65 70 75 80
 agc aca tac acc cag ctc tgg aac tgg gtt ccc gag tgc ttg agc tgt 288
 Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
 85 90 95
 ggc tcc cgc tgt agc tct gac cag gtg gaa act caa gcc tgc act cgg 336
 Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
 100 105 110

Sequence Listing

gaa cag aac cgc atc tgc acc tgc agg ccc ggc tgg tac tgc gcg ctg Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu 115 120 125	384
agc aag cag gag ggg tgc cgg ctg tgc gcg cgg ctg cgc aag tgc cgc Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg 130 135 140	432
ccg ggc ttc ggc gtg gcc aga cca gga act gaa aca tca gac gtg gtg Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val 145 150 155 160	480
tgc aag ccc tgt gcc cgg ggg acg ttc tcc aac acg act tca tcc acg Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr 165 170 175	528
gat att tgc agg ccc cac cag atc tgt aac gtg gtg gcc atc cct ggg Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly 180 185 190	576
aat gca agc atg gat gca gtc tgc acg tcc acg tcc ccc acc cgg agt Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser 195 200 205	624
atg gcc cca ggg gca gta cac tta ccc cag cca gtg tcc aca cga tcc Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser 210 215 220	672
caa cac acg cag cca act cca gaa ccc agc act gct cca agc acc tcc Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser 225 230 235 240	720
ttc ctg ctc cca atg ggc ccc agc ccc cca gct gaa ggg agc gga tcc Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Gly Ser 245 250 255	768
aac gca act aca ccc tac gcc cgg gag ccc ggg agc aca tgc cgg ctc Asn Ala Thr Thr Pro Tyr Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu 260 265 270	816

Sequence Listing

aga gaa tac tat gac cag aca gct cag atg tgc tgc agc aaa tgc tgc	864
Arg Glu Tyr Tyr Asp Gln Thr Ala Gln Met Cys Cys Ser Lys Cys Ser	
275 280 285	
ccg ggc caa cat gca aaa gtc ttc tgt acc aag acc tgc gac acc gtg	912
Pro Gly Gln His Ala Lys Val Phe Cys Thr Lys Thr Ser Asp Thr Val	
290 295 300	
tgt gac tcc tgt gag gac agc aca tac acc cag ctc tgg aac tgg gtt	960
Cys Asp Ser Cys Glu Asp Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val	
305 310 315 320	
ccc gag tgc ttg agc tgt ggc tcc cgc tgt agc tct gac cag gtg gaa	1008
Pro Glu Cys Leu Ser Cys Gly Ser Arg Cys Ser Ser Asp Gln Val Glu	
325 330 335	
act caa gcc tgc act cgg gaa cag aac cgc atc tgc acc tgc agg ccc	1056
Thr Gln Ala Cys Thr Arg Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro	
340 345 350	
ggc tgg tac tgc gcg ctg agc aag cag gag ggg tgc cgg ctg tgc gcg	1104
Gly Trp Tyr Cys Ala Leu Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala	
355 360 365	
ccg ctg cgc aag tgc cgc ccg ggc ttc ggc gtg gcc aga cca gga act	1152
Pro Leu Arg Lys Cys Arg Pro Gly Phe Gly Val Ala Arg Pro Gly Thr	
370 375 380	
gaa aca tca gac gtg gtg tgc aag ccc tgt gcc ccg ggg acg ttc tcc	1200
Glu Thr Ser Asp Val Val Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser	
385 390 395 400	
aac acg act tca tcc acg gat att tgc agg ccc cac cag atc tgt aac	1248
Asn Thr Thr Ser Ser Thr Asp Ile Cys Arg Pro His Gln Ile Cys Asn	
405 410 415	
gtg gtg gcc atc cct ggg aat gca agc atg gat gca gtc tgc acg tcc	1296
Val Val Ala Ile Pro Gly Asn Ala Ser Met Asp Ala Val Cys Thr Ser	
420 425 430	

Sequence Listing

acg tcc ccc acc cgg agt atg gcc cca ggg gca gta cac tta ccc cag	1344
Thr Ser Pro Thr Arg Ser Met Ala Pro Gly Ala Val His Leu Pro Gln	
435 440 445	
cca gtg tcc aca cga tcc caa cac acg cag cca act cca gaa ccc agc	1392
Pro Val Ser Thr Arg Ser Gln His Thr Gln Pro Thr Pro Glu Pro Ser	
450 455 460	
act gct cca agc acc tcc ttc ctg ctc cca atg ggc ccc agc ccc cca	1440
Thr Ala Pro Ser Thr Ser Phe Leu Leu Pro Met Gly Pro Ser Pro Pro	
465 470 475 480	
gct gaa ggg agc act ggc gac gca gag ccc aaa tct tgt gac aaa act	1488
Ala Glu Gly Ser Thr Gly Asp Ala Glu Pro Lys Ser Cys Asp Lys Thr	
485 490 495	
cac aca tgc cca ccg tgc cca gca cct gaa ctc ctg ggg gga ccg tca	1536
His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser	
500 505 510	
gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg atc tcc cgg	1584
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg	
515 520 525	
acc cct gag gtc aca tgc gtg gtg gtg gac gtg agc cac gaa gac cct	1632
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro	
530 535 540	
gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg cat aat gcc	1680
Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala	
545 550 555 560	
aag aca aag ccg cgg gag gag cag tac aac agc acg tac cgg gtg gtc	1728
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val	
565 570 575	
agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc aag gag tac	1776
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr	
580 585 590	

Sequence Listing

aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc gag aaa acc 1824
 Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
 595 600 605

atc tcc aaa gcc aaa ggg cag ccc cga gaa cca cag gtg tac acc ctg 1872
 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 610 615 620

ccc cca tcc cgg gat gag ctg acc aag aac cag gtc agc ctg acc tgc 1920
 Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
 625 630 635 640

ctg gtc aaa ggc ttc tat ccc agc gac atc gcc gtg gag tgg gag agc 1968
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 645 650 655

aat ggg cag cgg gag aac aac tac aag acc acg cct ccc gtg ctg gac 2016
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 660 665 670

tcc gac ggc tcc tcc ttc ctc tac agc aag ctc acc gtg gac aag agc 2064
 Ser Asp Gly Ser Ser Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 675 680 685

agg tgg cag cag ggg aac gtc ttc tca tgc tcc gtg atg cat gag gct 2112
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 690 695 700

ctg cac aac cac tac acg cag aag agc ctc tcc ctg tct ccg ggt aaa 2160
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 705 710 715 720

tga 2163

<210> 8

<211> 720

<212> PRT

<213> Homo sapiens

Sequence Listing

<400> 8

Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15

Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30

Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45

Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60

Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp
 65 70 75 80

Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys
 85 90 95

Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg
 100 105 110

Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu
 115 120 125

Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg
 130 135 140

Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val
 145 150 155 160

Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr
 165 170 175

Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly
 180 185 190

Asn Ala Ser Met Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser
 195 200 205

Sequence Listing

Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser
 210 215 220

Gln His Thr Gln Pro Thr Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser
 225 230 235 240

Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Gly Ser
 245 250 255

Asn Ala Thr Thr Pro Tyr Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu
 260 265 270

Arg Glu Tyr Tyr Asp Gln Thr Ala Gln Met Cys Cys Ser Lys Cys Ser
 275 280 285

Pro Gly Gln His Ala Lys Val Phe Cys Thr Lys Thr Ser Asp Thr Val
 290 295 300

Cys Asp Ser Cys Glu Asp Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val
 305 310 315 320

Pro Glu Cys Leu Ser Cys Gly Ser Arg Cys Ser Ser Asp Gln Val Glu
 325 330 335

Thr Gln Ala Cys Thr Arg Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro
 340 345 350

Gly Trp Tyr Cys Ala Leu Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala
 355 360 365

Pro Leu Arg Lys Cys Arg Pro Gly Phe Gly Val Ala Arg Pro Gly Thr
 370 375 380

Glu Thr Ser Asp Val Val Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser
 385 390 395 400

Asn Thr Thr Ser Ser Thr Asp Ile Cys Arg Pro His Gln Ile Cys Asn
 405 410 415

Val Val Ala Ile Pro Gly Asn Ala Ser Met Asp Ala Val Cys Thr Ser

Sequence Listing

420	425	430
Thr Ser Pro Thr Arg Ser Met Ala Pro Gly Ala Val His Leu Pro Gln		
435	440	445
Pro Val Ser Thr Arg Ser Gln His Thr Gln Pro Thr Pro Glu Pro Ser		
450	455	460
Thr Ala Pro Ser Thr Ser Phe Leu Leu Pro Met Gly Pro Ser Pro Pro		
465	470	475 480
Ala Glu Gly Ser Thr Gly Asp Ala Glu Pro Lys Ser Cys Asp Lys Thr		
485	490	495
His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser		
500	505	510
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg		
515	520	525
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro		
530	535	540
Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala		
545	550	555 560
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val		
565	570	575
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr		
580	585	590
Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr		
595	600	605
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu		
610	615	620
Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys		
625	630	635 640

Sequence Listing

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 645 650 655

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 660 665 670

Ser Asp Gly Ser Ser Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 675 680 685

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 690 695 700

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 705 710 715 720

<210> 9
 <211> 1827
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1824)
 <223> mgTNFR1-TNFR1-IgG

<220>
 <221> C_region
 <222> (1126)..(1827)
 <223> Hinge, CH2, CH3 region

<220>
 <221> misc_signal
 <222> (160)..(168)
 <223> N-linked glycosylation site

Sequence Listing

<220>
<221> misc_signal
<222> (433)..(441)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (451)..(459)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (565)..(573)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (574)..(582)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (592)..(600)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (610)..(618)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (925)..(933)
<223> N-linked glycosylation site

Sequence Listing

<220>
<221> misc_signal
<222> (943)..(951)
<223> N-linked glycosylation site

<220>
<221> primer_bind
<222> (1)..(15)
<223> PCR primer SEQ ID : 25 binding site

<220>
<221> primer_bind
<222> (545)..(606)
<223> PCR primer SEQ ID : 37(antisense) binding site

<220>
<221> primer_bind
<222> (559)..(621)
<223> PCR primer SEQ ID : 36 binding site

<220>
<221> primer_bind
<222> (1108)..(1144)
<223> PCR primer SEQ ID : 26(antisense) binding site

<220>
<221> primer_bind
<222> (1108)..(1144)
<223> PCR primer SEQ ID : 27 binding site

<220>
<221> primer_bind

Sequence Listing

<222> (1804)..(1827)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(60)

<223> signal peptide

<400> 9

atg ggc ctc tcc acc gtg cct gac ctg ctg cgc ctg gtg ctc ctg	48
Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu	
1 5 10 15	

gag ctg ttg gtg gga ata tac ccc tca ggg gtt att gga ctg gtc cct	96
Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro	
20 25 30	

cac cta ggg gac agg gag aag aga gat agt gtg tgt ccc caa gga aaa	144
His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys	
35 40 45	

tat atc cac cct caa aat aat tgc att tgc tgt acc aag tgc cac aaa	192
Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys	
50 55 60	

gga acc tac ttg tac aat gac tgt cca ggc cgc ggg cag gat acg gac	240
Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp	
65 70 75 80	

tgc agg gag tgt gag agc ggc tcc ttc acc gct tca gaa aac cac ctc	288
Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu	
85 90 95	

aga cac tgc ctc agc tgc tcc aaa tgc cga aag gaa atg ggt cag gtg	336
Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val	
100 105 110	

gag atc tct tct tgc aca gtg gac cgc gac acc gtg tgt ggc tgc agg	384
---	-----

Sequence Listing

Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg	
115	120
125	
aag aac cag tac cgg cat tat tgg agt gaa aac ctt ttc cag tgc ttc	432
Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe	
130	135
140	
aat tgc agc ctc tgc ctc aat ggg acc gtg cac ctc tcc tgc cag gag	480
Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu	
145	150
155	160
aaa cag aac acc gtg tgc acc tgc cat gca ggt ttc ttt cta aga gaa	528
Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu	
165	170
175	
aac gag tgt gtc tcc tgt agt aac tgt aag aaa agc aac gag acc aac	576
Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Asn Glu Thr Asn	
180	185
190	
aag acc tgc cta cac aac ggg tcc agg gag aag aac gat agt gtg tgt	624
Lys Thr Cys Leu His Asn Gly Ser Arg Glu Lys Asn Asp Ser Val Cys	
195	200
205	
ccc caa gga aaa tat atc cac cct caa aat aat tgg att tgc tgt acc	672
Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr	
210	215
220	
aag tgc cac aaa gga acc tac ttg tac aat gac tgt cca ggc ccg ggg	720
Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly	
225	230
235	240
cag gat acg gac tgc agg gag tgt gag agc ggc tcc ttc acc gct tca	768
Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser	
245	250
255	
gaa aac cac ctc aga cac tgc ctc agc tgc tcc aaa tgc cga aag gaa	816
Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu	
260	265
270	
atg ggt cag gtg gag atc tct tct tgc aca gtg gac cgg gac acc gtg	864

Sequence Listing

Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val	
275	280 285
tgt ggc tgc agg aag aac cag tac cgg cat tat tgg agt gaa aac ctt	912
Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu	
290	295 300
ttc cag tgc ttc aat tgc agc ctc tgc ctc aat ggg acc gtg cac ctc	960
Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu	
305	310 315 320
tcc tgc cag gag aaa cag aac acc gtg tgc acc tgc cat gca ggt ttc	1008
Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe	
325	330 335
ttt cta aga gaa aac gag tgt gtc tcc tgt agt aac tgt aag aaa agc	1056
Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser	
340	345 350
ctg gag tgc acg aag ttg tgc cta ccc cag att gag aat gtt aag ggc	1104
Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly	
355	360 365
act gag gac tca ggc acc aca gca gag ccc aaa tct tgt gac aaa act	1152
Thr Glu Asp Ser Gly Thr Thr Ala Glu Pro Lys Ser Cys Asp Lys Thr	
370	375 380
cac aca tgc cca ccg tgc cca gca cct gaa ctc ctg ggg gga ccg tca	1200
His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser	
385	390 395 400
gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg atc tcc cgg	1248
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg	
405	410 415
acc cct gag gtc aca tgc gtg gtg gtg gac gtg agc cac gaa gac cct	1296
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro	
420	425 430
gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg cat aat gcc	1344

Sequence Listing

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala	
435	440
	445
aag aca aag ccg cgg gag gag cag tac aac agc acg tac cgg gtg gtc	1392
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val	
450	455
	460
agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc aag gag tac	1440
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr	
465	470
	475
	480
aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc gag aaa acc	1488
Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr	
485	490
	495
atc tcc aaa gcc aaa ggg cag ccc cga gaa cca cag gtg tac acc ctg	1536
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu	
500	505
	510
ccc cca tcc cgg gat gag ctg acc aag aac cag gtc agc ctg acc tgc	1584
Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys	
515	520
	525
ctg gtc aaa ggc ttc tat ccc agc gac atc gcc gtg gag tgg gag agc	1632
Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser	
530	535
	540
aat ggg cag ccg gag aac aac tac aag acc acg cct ccc gtg ctg gac	1680
Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp	
545	550
	555
	560
tcc gac ggc tcc ttc ttc ctc tac agc aag ctc acc gtg gac aag agc	1728
Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser	
565	570
	575
agg tgg cag cag ggg aac gtc ttc tca tgc tcc gtg atg cat gag gct	1776
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala	
580	585
	590
ctg cac aac cac tac acg cag aag agc ctc tcc ctg tct ccg ggt aaa	1824

Sequence Listing

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 595 600 605

tga

1827

<210> 10

<211> 608

<212> PRT

<213> Homo sapiens

<400> 10

Met Gly Leu Ser Thr Val Pro Asp Leu Leu Leu Pro Leu Val Leu Leu
 1 5 10 15

Glu Leu Leu Val Gly Ile Tyr Pro Ser Gly Val Ile Gly Leu Val Pro
 20 25 30

His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro Gln Gly Lys
 35 40 45

Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys
 50 55 60

Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp
 65 70 75 80

Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu
 85 90 95

Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val
 100 105 110

Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg
 115 120 125

Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe
 130 135 140

Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu

Sequence Listing

145	150	155	160
Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu			
165	170	175	
Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Asn Glu Thr Asn			
180	185	190	
Lys Thr Cys Leu His Asn Gly Ser Arg Glu Lys Asn Asp Ser Val Cys			
195	200	205	
Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr			
210	215	220	
Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly			
225	230	235	240
Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser			
245	250	255	
Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu			
260	265	270	
Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val			
275	280	285	
Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu			
290	295	300	
Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu			
305	310	315	320
Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe			
325	330	335	
Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser			
340	345	350	
Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly			
355	360	365	

Sequence Listing

Thr Glu Asp Ser Gly Thr Thr Ala Glu Pro Lys Ser Cys Asp Lys Thr
 370 375 380

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
 385 390 395 400

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 405 410 415

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
 420 425 430

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 435 440 445

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val
 450 455 460

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 465 470 475 480

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
 485 490 495

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 500 505 510

Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
 515 520 525

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 530 535 540

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 545 550 555 560

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 565 570 575

Sequence Listing

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 580 585 590

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 595 600 605

<210> 11
 <211> 1980
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1977)
 <223> mgTNER2-TNFR2-IgG

<220>
 <221> C_region
 <222> (1279)..(1980)
 <223> Hinge, CH2, CH3 region

<220>
 <221> misc_signal
 <222> (511)..(519)
 <223> N-linked glycosylation site

<220>
 <221> misc_signal
 <222> (577)..(585)
 <223> N-linked glycosylation site

<220>
 <221> misc_signal

Sequence Listing

<222> (595)..(603)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (616)..(624)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (1018)..(1026)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (1084)..(1092)

<223> N-linked glycosylation site

<220>

<221> primer_bind

<222> (1)..(15)

<223> PCR primer SEQ ID : 29 binding site

<220>

<221> primer_bind

<222> (586)..(627)

<223> PCR primer SEQ ID : 39(antisense) binding site

<220>

<221> primer_bind

<222> (586)..(630)

<223> PCR primer SEQ ID : 38 binding site

Sequence Listing

<220>
 <221> primer_bind
 <222> (1261)..(1296)
 <223> PCR primer SEQ ID : 30(antisense) binding site

<220>
 <221> primer_bind
 <222> (1261)..(1296)
 <223> PCR primer SEQ ID : 31 binding site

<220>
 <221> primer_bind
 <222> (1957)..(1980)
 <223> PCR primer SEQ ID : 28(antisense) binding site

<220>
 <221> sig_peptide
 <222> (1)..(66)
 <223> signal peptide

<400> 11
 atg gcg ccc gtc gcc gtc tgg gcc gcg ctg gcc gtc gga ctg gag ctc 48
 Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu
 1 5 10 15
 tgg gct gcg gcg cac gcc ttg ccc gcc cag gtg gca ttt aca ccc tac 96
 Trp Ala Ala Ala His Ala Leu Pro Ala Gln Val Ala Phe Thr Pro Tyr
 20 25 30
 gcc ccg gag ccc ggg agc aca tgc cgg ctc aga gaa tac tat gac cag 144
 Ala Pro Glu Pro Gly Ser Thr Cys Arg Leu Arg Glu Tyr Tyr Asp Gln
 35 40 45
 aca gct cag atg tgc tgc agc aaa tgc tcg ccg ggc caa cat gca aaa 192
 Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys
 50 55 60

- 48 -

Sequence Listing

aaa tgc tgc ccg ggc caa cat gca aaa gtc ttc tgt acc aag acc tgc	720
Lys Cys Ser Pro Gly Gln His Ala Lys Val Phe Cys Thr Lys Thr Ser	
225 230 235 240	
gac acc gtg tgt gac tcc tgt gag gac agc aca tac acc cag ctc tgg	768
Asp Thr Val Cys Asp Ser Cys Glu Asp Ser Thr Tyr Thr Gln Leu Trp	
245 250 255	
aac tgg gtt ccc gag tgc ttg agc tgt ggc tcc cgc tgt agc tct gac	816
Asn Trp Val Pro Glu Cys Leu Ser Cys Gly Ser Arg Cys Ser Ser Asp	
260 265 270	
cag gtg gaa act caa gcc tgc act cgg gaa cag aac cgc atc tgc acc	864
Gln Val Glu Thr Gln Ala Cys Thr Arg Glu Gln Asn Arg Ile Cys Thr	
275 280 285	
tgc agg ccc ggc tgg tac tgc gcg ctg agc aag cag gag ggg tgc cgg	912
Cys Arg Pro Gly Trp Tyr Cys Ala Leu Ser Lys Gln Glu Gly Cys Arg	
290 295 300	
ctg tgc gcg ccg ctg cgc aag tgc cgc ccg ggc ttc gcc gtg gcc aga	960
Leu Cys Ala Pro Leu Arg Lys Cys Arg Pro Gly Phe Gly Val Ala Arg	
305 310 315 320	
cca gga act gaa aca tca gac gtg gtg tgc aag ccc tgt gcc ccg ggg	1008
Pro Gly Thr Glu Thr Ser Asp Val Val Cys Lys Pro Cys Ala Pro Gly	
325 330 335	
acg ttc tcc aac acg act tca tcc acg gat att tgc agg ccc cac cag	1056
Thr Phe Ser Asn Thr Thr Ser Ser Thr Asp Ile Cys Arg Pro His Gln	
340 345 350	
atc tgt aac gtg gtg gcc atc cct ggg aat gca agc atg gat gca gtc	1104
Ile Cys Asn Val Val Ala Ile Pro Gly Asn Ala Ser Met Asp Ala Val	
355 360 365	
tgc acg tcc acg tcc ccc acc cgg agt atg gcc cca ggg gca gta cac	1152
Cys Thr Ser Thr Ser Pro Thr Arg Ser Met Ala Pro Gly Ala Val His	
370 375 380	

Sequence Listing

tta ccc cag cca gtg tcc aca cga tcc caa cac acg cag cca act cca	1200
Leu Pro Gln Pro Val Ser Thr Arg Ser Gln His Thr Gln Pro Thr Pro	
385 390 395 400	
gaa ccc agc act gct cca agc acc tcc ttc ctg ctc cca atg ggc ccc	1248
Glu Pro Ser Thr Ala Pro Ser Thr Ser Phe Leu Leu Pro Met Gly Pro	
405 410 415	
agc ccc cca gct gaa ggg agc act ggc gac gca gag ccc aaa tct tgt	1296
Ser Pro Pro Ala Glu Gly Ser Thr Gly Asp Ala Glu Pro Lys Ser Cys	
420 425 430	
gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa ctc ctg ggg	1344
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly	
435 440 445	
gga ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg	1392
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met	
450 455 460	
atc tcc cgg acc cct gag gtc aca tgc gtg gtg gac gtg agc cac	1440
Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His	
465 470 475 480	
gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg	1488
Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val	
485 490 495	
cat aat gcc aag aca aag ccg cgg gag gag cag tac aac agc acg tac	1536
His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr	
500 505 510	
cgg gtg gtc agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc	1584
Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly	
515 520 525	
aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc	1632
Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile	
530 535 540	

Sequence Listing

```

gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa cca cag gtg      1680
Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
545                      550                      555                      560

tac acc ctg ccc cca tcc cgg gat gag ctg acc aag aac cag gtc agc      1728
Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser
                    565                      570                      575

ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc gcc gtg gag      1776
Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
                    580                      585                      590

tgg gag agc aat ggg cag ccg gag aac aac tac aag acc acg cct ccc      1824
Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
                    595                      600                      605

gtg ctg gac tcc gac ggc tcc ttc ttc ctc tac agc aag ctc acc gtg      1872
Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
                    610                      615                      620

gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc tcc gtg atg      1920
Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
625                      630                      635                      640

cat gag gct ctg cac aac cac tac acg cag aag agc ctc tcc ctg tct      1968
His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
                    645                      650                      655

ccg ggt aaa          tga          1980
Pro Gly Lys

```

```

<210> 12
<211> 659
<212> PRT
<213> Homo sapiens

```

```

<400> 12
Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu

```

Sequence Listing

1	5	10	15
Trp	Ala	Ala	Ala
His	Ala	Leu	Pro
Ala	Gln	Val	Ala
Phe	Thr	Pro	Tyr
20	25	30	
Ala	Pro	Glu	Pro
Gly	Ser	Thr	Cys
Arg	Leu	Arg	Glu
Tyr	Tyr	Asp	Gln
35	40	45	
Thr	Ala	Gln	Met
Cys	Cys	Ser	Lys
Cys	Ser	Pro	Gly
Gln	His	Ala	Lys
50	55	60	
Val	Phe	Cys	Thr
Lys	Thr	Ser	Asp
Thr	Val	Cys	Asp
Ser	Cys	Glu	Asp
65	70	75	80
Ser	Thr	Tyr	Thr
Gln	Leu	Trp	Asn
Trp	Val	Pro	Glu
Cys	Leu	Ser	Cys
85	90	95	
Gly	Ser	Arg	Cys
Ser	Ser	Asp	Gln
Val	Glu	Thr	Gln
Ala	Cys	Thr	Arg
100	105	110	
Glu	Gln	Asn	Arg
Ile	Cys	Thr	Cys
Arg	Pro	Gly	Trp
Tyr	Cys	Ala	Leu
115	120	125	
Ser	Lys	Gln	Glu
Gly	Cys	Arg	Leu
Cys	Ala	Pro	Leu
Arg	Lys	Cys	Arg
130	135	140	
Pro	Gly	Phe	Gly
Val	Ala	Arg	Pro
Gly	Thr	Glu	Thr
Ser	Asp	Val	Val
145	150	155	160
Cys	Lys	Pro	Cys
Ala	Pro	Gly	Thr
Phe	Ser	Asn	Thr
Thr	Thr	Ser	Ser
Thr	165	170	175
Asp	Ile	Cys	Arg
Pro	His	Gln	Ile
Cys	Asn	Val	Val
Ala	Ile	Pro	Gly
180	185	190	
Asn	Ala	Ser	Met
Asp	Ala	Asn	Cys
Thr	Ser	Pro	Glu
Pro	Asn	Ser	Thr
195	200	205	
Cys	Arg	Leu	Arg
Glu	Tyr	Tyr	Asp
Gln	Thr	Ala	Gln
Met	Cys	Cys	Ser
210	215	220	

Sequence Listing

Lys Cys Ser Pro Gly Gln His Ala Lys Val Phe Cys Thr Lys Thr Ser
 225 230 235 240

Asp Thr Val Cys Asp Ser Cys Glu Asp Ser Thr Tyr Thr Gln Leu Trp
 245 250 255

Asn Trp Val Pro Glu Cys Leu Ser Cys Gly Ser Arg Cys Ser Ser Asp
 260 265 270

Gln Val Glu Thr Gln Ala Cys Thr Arg Glu Gln Asn Arg Ile Cys Thr
 275 280 285

Cys Arg Pro Gly Trp Tyr Cys Ala Leu Ser Lys Gln Glu Gly Cys Arg
 290 295 300

Leu Cys Ala Pro Leu Arg Lys Cys Arg Pro Gly Phe Gly Val Ala Arg
 305 310 315 320

Pro Gly Thr Glu Thr Ser Asp Val Val Cys Lys Pro Cys Ala Pro Gly
 325 330 335

Thr Phe Ser Asn Thr Thr Ser Ser Thr Asp Ile Cys Arg Pro His Gln
 340 345 350

Ile Cys Asn Val Val Ala Ile Pro Gly Asn Ala Ser Met Asp Ala Val
 355 360 365

Cys Thr Ser Thr Ser Pro Thr Arg Ser Met Ala Pro Gly Ala Val His
 370 375 380

Leu Pro Gln Pro Val Ser Thr Arg Ser Gln His Thr Gln Pro Thr Pro
 385 390 395 400

Glu Pro Ser Thr Ala Pro Ser Thr Ser Phe Leu Leu Pro Met Gly Pro
 405 410 415

Ser Pro Pro Ala Glu Gly Ser Thr Gly Asp Ala Glu Pro Lys Ser Cys
 420 425 430

Sequence Listing

```

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
    435                440                445

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
    450                455                460

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
    465                470                475                480

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
                485                490                495

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
                500                505                510

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
    515                520                525

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
    530                535                540

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
    545                550                555                560

Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser
                565                570                575

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
    580                585                590

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
    595                600                605

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
    610                615                620

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
    625                630                635                640

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser

```

Sequence Listing

645

650

655

Pro Gly Lys

<210> 13
<211> 1314
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(1311)
<223> CD2-IgG

<220>
<221> C_region
<222> (613)..(1314)
<223> Hinge, CH2, CH3 region

<220>
<221> misc_signal
<222> (265)..(273)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (421)..(429)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (448)..(456)
<223> N-linked glycosylation site

Sequence Listing

<220>
 <221> primer_bind
 <222> (1)..(27)
 <223> PCR primer SEQ ID : 40 binding site

<220>
 <221> primer_bind
 <222> (589)..(618)
 <223> PCR primer SEQ ID : 41(antisense) binding site

<220>
 <221> primer_bind
 <222> (611)..(633)
 <223> PCR primer SEQ ID : 42 binding site

<220>
 <221> primer_bind
 <222> (1292)..(1314)
 <223> PCR primer SEQ ID : 28(antisense) binding site

<220>
 <221> sig_peptide
 <222> (1)..(72)
 <223> signal peptide

<400> 13
 atg agc ttt cca tgt aaa ttt gta gcc agc ttc ctt ctg att ttc aat 48
 Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn
 1 5 10 15
 gtt tct tcc aaa ggt gca gtc tcc aaa gag att acg aat gcc ttg gaa 96
 Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu
 20 25 30

Sequence Listing

acc tgg ggt gcc ttg ggt cag gac atc aac ttg gac att cct agt ttt	144
Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe	
35 40 45	
caa atg agt gat gat att gac gat ata aaa tgg gaa aaa act tca gac	192
Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp	
50 55 60	
aag aaa aag att gca caa ttc aga aaa gag aaa gag act ttc aag gaa	240
Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu	
65 70 75 80	
aaa gat aca tat aag cta ttt aaa aat gga act ctg aaa att aag cat	288
Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His	
85 90 95	
ctg aag acc gat gat cag gat atc tac aag gta tca ata tat gat aca	336
Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr	
100 105 110	
aaa gga aaa aat gtg ttg gaa aaa ata ttt gat ttg aag att caa gag	384
Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu	
115 120 125	
agg gtc tca aaa cca aag atc tcc tgg act tgt atc aac aca acc ctg	432
Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu	
130 135 140	
acc tgt gag gta atg aat gga act gac ccc gaa tta aac ctg tat caa	480
Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln	
145 150 155 160	
gat ggg aaa cat cta aaa ctt tct cag agg gtc atc aca cac aag tgg	528
Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp	
165 170 175	
acc acc agc ctg agt gca aaa ttc aag tgc aca gca ggg aac aaa gtc	576
Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val	
180 185 190	

Sequence Listing

agc aag gaa tcc agt gtc gag cct gtc agc tgt cct gca gag ccc aaa	624
Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro Ala Glu Pro Lys	
195 200 205	
tct tgt gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa ctc	672
Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu	
210 215 220	
ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac acc	720
Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr	
225 230 235 240	
ctc atg atc tcc ccg acc cct gag gtc aca tgc gtg gtg gac gtg	768
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val	
245 250 255	
agc cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc gtg	816
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val	
260 265 270	
gag gtg cat aat gcc aag aca aag ccg ccg gag gag cag tac aac agc	864
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser	
275 280 285	
acg tac ccg gtg gtc agc gtc ctc acc gtc ctg cac cag gac tgg ctg	912
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu	
290 295 300	
aat ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca gcc	960
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala	
305 310 315 320	
ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa cca	1008
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro	
325 330 335	
cag gtg tac acc ctg ccc cca tcc ccg gat gag ctg acc aag aac cag	1056
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln	
340 345 350	

Sequence Listing

gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc gcc 1104
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 355 360 365

gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag acc acg 1152
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 370 375 380

cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tac agc aag ctc 1200
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 385 390 395 400

acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc tcc 1248
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 405 410 415

gtg atg cat gag gct ctg cac aac cac tac acg cag aag agc ctc tcc 1296
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 420 425 430

ctg tct ccg ggt aaa tga 1314
 Leu Ser Pro Gly Lys
 435

<210> 14
 <211> 437
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn
 1 5 10 15

Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu
 20 25 30

Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe
 35 40 45

Sequence Listing

Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp
50 55 60

Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu
65 70 75 80

Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His
85 90 95

Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr
100 105 110

Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu
115 120 125

Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu
130 135 140

Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln
145 150 155 160

Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp
165 170 175

Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val
180 185 190

Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro Ala Glu Pro Lys
195 200 205

Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu
210 215 220

Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
225 230 235 240

Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
245 250 255

Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val

Sequence Listing

260	265	270
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser		
275	280	285
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu		
290	295	300
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala		
305	310	315 320
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro		
325	330	335
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln		
340	345	350
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala		
355	360	365
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr		
370	375	380
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu		
385	390	395 400
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser		
405	410	415
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser		
420	425	430
Leu Ser Pro Gly Lys		
435		

<210> 15
 <211> 1134
 <212> DNA
 <213> Homo sapiens

Sequence Listing

<220>

<221> CDS

<222> (1)..(1131)

<223> CTLA4-IgG

<220>

<221> C_region

<222> (433)..(1134)

<223> Hinge, CH2, CH3 region

<220>

<221> misc_signal

<222> (289)..(297)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (385)..(393)

<223> N-linked glycosylation site

<220>

<221> primer_bind

<222> (1)..(15)

<223> PCR primer SEQ ID : 43 binding site

<220>

<221> primer_bind

<222> (409)..(438)

<223> PCR primer SEQ ID : 44(antisense) binding site

<220>

<221> primer_bind

<222> (430)..(453)

Sequence Listing

<223> PCR primer SEQ ID : 42 binding site

<220>

<221> primer_bind

<222> (1111)..(1134)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(63)

<223> signal peptide

<400> 15

atg agg acc tgg ccc tgc act ctc ctg ttt ttt ctt ctc ttc atc cct 48

Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro

1 5 10 15

gtc ttc tgc aaa gca atg cac gtg gcc cag cct gct gtg gta ctg gcc 96

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala

20 25 30

agc agc cga ggc atc gcc agc ttt gtg tgt gag tat gca tct cca gcc 144

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly

35 40 45

aaa gcc act gag gtc cgg gtg aca gtg ctt cgg cag gct gac agc cag 192

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln

50 55 60

gtg act gaa gtc tgt gcg gca acc tac atg atg ggg aat gag ttg acc 240

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr

65 70 75 80

ttc cta gat gat tcc atc tgc acg ggc acc tcc agt gga aat caa gtg 288

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val

85 90 95

Sequence Listing

aac ctc act atc caa gga ctg agg gcc atg gac acg gga ctc tac atc	336
Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile	
100 105 110	
tgc aag gtg gag ctc atg tac cca ccg cca tac tac ctg ggc ata ggc	384
Cys Lys Val Glu Leu Met Tyr Pro Pro Tyr Tyr Leu Gly Ile Gly	
115 120 125	
aac gga acc cag att tat gta att gat cca gaa ccg tgc cca gat tct	432
Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser	
130 135 140	
gca gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg tgc cca	480
Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro	
145 150 155 160	
gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa	528
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys	
165 170 175	
ccc aag gac acc ctc atg atc tcc ccg acc cct gag gtc aca tgc gtg	576
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val	
180 185 190	
gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac	624
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr	
195 200 205	
gtg gac ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag	672
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu	
210 215 220	
cag tac aac agc acg tac ccg gtg gtc agc gtc ctc acc gtc ctg cac	720
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His	
225 230 235 240	
cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa	768
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys	
245 250 255	

Sequence Listing

```

gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag      816
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
      260              265              270

ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat gag ctg      864
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
      275              280              285

acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc      912
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
      290              295              300

agc gac atc gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac      960
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
305              310              315              320

tac aag acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc      1008
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
      325              330              335

tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc      1056
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
      340              345              350

ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg cag      1104
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
      355              360              365

aag agc ctc tcc ctg tct ccg ggt aaa      tga      1134
Lys Ser Leu Ser Leu Ser Pro Gly Lys
      370              375

```

<210> 16

<211> 377

<212> PRT

<213> Homo sapiens

<400> 16

Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro

Sequence Listing

```

1           5           10          15
Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
      20           25           30
Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
      35           40           45
Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
      50           55           60
Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
      65           70           75           80
Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
      85           90           95
Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
      100          105          110
Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
      115          120          125
Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser
      130          135          140
Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
      145          150          155          160
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
      165          170          175
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
      180          185          190
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
      195          200          205
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
      210          215          220

```

Sequence Listing

Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
225 230 235 240

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
245 250 255

Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
260 265 270

Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
275 280 285

Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
290 295 300

Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
305 310 315 320

Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
325 330 335

Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
340 345 350

Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
355 360 365

Lys Ser Leu Ser Leu Ser Pro Gly Lys
370 375

<210> 17
<211> 1854
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(1851)

Sequence Listing

<223> CD2-CD2-IgG

<220>

<221> C_region

<222> (1153)..(1854)

<223> Hinge, CH2, CH3 region

<220>

<221> misc_signal

<222> (265)..(273)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (421)..(429)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (448)..(456)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (805)..(813)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (961)..(969)

<223> N-linked glycosylation site

<220>

Sequence Listing

<221> misc_signal
<222> (988)..(996)
<223> N-linked glycosylation site

<220>
<221> primer_bind
<222> (1)..(27)
<223> PCR primer SEQ ID : 40 binding site

<220>
<221> primer_bind
<222> (598)..(612)
<223> PCR primer SEQ ID : 46(antisense) binding site

<220>
<221> primer_bind
<222> (612)..(630)
<223> PCR primer SEQ ID : 45 binding site

<220>
<221> primer_bind
<222> (1128)..(1158)
<223> PCR primer SEQ ID : 41(antisense) binding site

<220>
<221> primer_bind
<222> (1151)..(1173)
<223> PCR primer SEQ ID : 42 binding site

<220>
<221> primer_bind
<222> (1832)..(1854)
<223> PCR primer SEQ ID : 28(antisense) binding site

Sequence Listing

<220>

<221> sig_peptide

<222> (1)..(72)

<223> signal peptide

<400> 17

atg agc ttt cca tgt aaa ttt gta gcc agc ttc ctt ctg att ttc aat 48

Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn

1 5 10 15

gtt tct tcc aaa ggt gca gtc tcc aaa gag att acg aat gcc ttg gaa 96

Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu

20 25 30

acc tgg ggt gcc ttg ggt cag gac atc aac ttg gac att cct agt ttt 144

Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe

35 40 45

caa atg agt gat gat att gac gat ata aaa tgg gaa aaa act tca gac 192

Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp

50 55 60

aag aaa aag att gca caa ttc aga aaa gag aaa gag act ttc aag gaa 240

Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu

65 70 75 80

aaa gat aca tat aag cta ttt aaa aat gga act ctg aaa att aag cat 288

Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His

85 90 95

ctg aag acc gat gat cag gat atc tac aag gta tca ata tat gat aca 336

Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr

100 105 110

aaa gga aaa aat gty ttg gaa aaa ata ttt gat ttg aag att caa gag 384

Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu

115 120 125

Sequence Listing

agg gtc tca aaa cca aag atc tcc tgg act tgt atc aac aca acc ctg	432
Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu	
130 135 140	
acc tgt gag gta atg aat gga act gac ccc gaa tta aac ctg tat caa	480
Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln	
145 150 155 160	
gat ggg aaa cat cta aaa ctt tct cag agy gtc atc aca cac aag tgg	528
Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp	
165 170 175	
acc acc agc ctg agt gca aaa ttc aag tgc aca gca ggg aac aaa gtc	576
Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val	
180 185 190	
agc aag gaa tcc agt gtc gag cct gtc agc tgt cct aaa gag att acg	624
Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro Lys Glu Ile Thr	
195 200 205	
aat gcc ttg gaa acc tgg ggt gcc ttg ggt cag gac atc aac ttg gac	672
Asn Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp	
210 215 220	
att cct agt ttt caa atg agt gat gat att gac gat ata aaa tgg gaa	720
Ile Pro Ser Phe Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu	
225 230 235 240	
aaa act tca gac aag aaa aag att gca caa ttc aga aaa gag aaa gag	768
Lys Thr Ser Asp Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu	
245 250 255	
act ttc aag gaa aaa gat aca tat aag cta ttt aaa aat gga act ctg	816
Thr Phe Lys Glu Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu	
260 265 270	
aaa att aag cat ctg aag acc gat gat cag gat atc tac aag gta tca	864
Lys Ile Lys His Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser	
275 280 285	

Sequence Listing

ata tat gat aca aaa gga aaa aat gtg ttg gaa aaa ata ttt gat ttg	912
Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu	
290 295 300	
aag att caa gag agg gtc tca aaa cca aag atc tcc tgg act tgt atc	960
Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile	
305 310 315 320	
aac aca acc ctg acc tgt gag gta atg aat gga act gac ccc gaa tta	1008
Asn Thr Thr Leu Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu	
325 330 335	
aac ctg tat caa gat ggg aaa cat cta aaa ctt tct cag agg gtc atc	1056
Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile	
340 345 350	
aca cac aag tgg acc acc agc ctg agt gca aaa ttc aag tgc aca gca	1104
Thr His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala	
355 360 365	
ggg aac aaa gtc agc aag gaa tcc agt gtc gag cct gtc agc tgt cct	1152
Gly Asn Lys Val Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro	
370 375 380	
gca gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg tgc cca	1200
Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro	
385 390 395 400	
gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa	1248
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys	
405 410 415	
ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg	1296
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val	
420 425 430	
gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac	1344
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr	
435 440 445	

Sequence Listing

gtg gac ggc gtg gag gtg cat aat gcc aag aca aag cgg cgg gag gag Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu 450 455 460	1392
cag tac aac agc acg tac cgg gtg gtc agc gtc ctc acc gtc tgt cac Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His 465 470 475 480	1440
cag gac tgg ctg aat gcc aag gag tac aag tgc aag gtc tcc aac aaa Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys 485 490 495	1488
gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln 500 505 510	1536
ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat gag ctg Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu 515 520 525	1584
acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa gcc ttc tat ccc Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro 530 535 540	1632
agc gac atc gcc gtg gag tgg gag agc aat ggg cag cgg gag aac aac Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn 545 550 555 560	1680
tac aag acc acg cct ccc gtg ctg gac tcc gac gcc tcc ttc ttc ctc Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu 565 570 575	1728
tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val 580 585 590	1776
ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg cag Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln 595 600 605	1824

Sequence Listing

aag agc ctc tcc ctg tct ccg ggt aaa tga 1854
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 610 615

<210> 18
 <211> 617
 <212> PRT
 <213> Homo sapiens

<400> 18
 Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn
 1 5 10 15

Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu
 20 25 30

Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe
 35 40 45

Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp
 50 55 60

Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu
 65 70 75 80

Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His
 85 90 95

Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr
 100 105 110

Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu
 115 120 125

Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu
 130 135 140

Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln
 145 150 155 160

Sequence Listing

Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp
165 170 175

Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val
180 185 190

Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro Lys Glu Ile Thr
195 200 205

Asn Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp
210 215 220

Ile Pro Ser Phe Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu
225 230 235 240

Lys Thr Ser Asp Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu
245 250 255

Thr Phe Lys Glu Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu
260 265 270

Lys Ile Lys His Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser
275 280 285

Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu
290 295 300

Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile
305 310 315 320

Asn Thr Thr Leu Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu
325 330 335

Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile
340 345 350

Thr His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala
355 360 365

Sequence Listing

Gly Asn Lys Val Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro
370 375 380

Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
385 390 395 400

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
405 410 415

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
420 425 430

Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
435 440 445

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
450 455 460

Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His
465 470 475 480

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
485 490 495

Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
500 505 510

Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
515 520 525

Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
530 535 540

Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
545 550 555 560

Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
565 570 575

Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val

Sequence Listing

580	585	590
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln		
595	600	605
Lys Ser Leu Ser Leu Ser Pro Gly Lys		
610	615	

<210> 19
 <211> 1509
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1506)
 <223> CTLA4-CTLA4-IgG

<220>
 <221> C_region
 <222> (808)..(1509)
 <223> Hinge, CH2, CH3 region

<220>
 <221> misc_signal
 <222> (289)..(297)
 <223> N-linked glycosylation site

<220>
 <221> misc_signal
 <222> (385)..(393)
 <223> N-linked glycosylation site

<220>
 <221> misc_signal

Sequence Listing

<222> (664)..(672)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (760)..(768)
<223> N-linked glycosylation site

<220>
<221> primer_bind
<222> (1)..(15)
<223> PCR primer SEQ ID : 43 binding site

<220>
<221> primer_bind
<222> (418)..(431)
<223> PCR primer SEQ ID : 48(antisense) binding site

<220>
<221> primer_bind
<222> (432)..(453)
<223> PCR primer SEQ ID : 47 binding site

<220>
<221> primer_bind
<222> (784)..(813)
<223> PCR primer SEQ ID : 44(antisense) binding site

<220>
<221> primer_bind
<222> (805)..(826)
<223> PCR primer SEQ ID : 42 binding site

Sequence Listing

<220>

<221> primer_bind

<222> (1486)..(1509)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(63)

<223> signal peptide

<400> 19

atg agg acc tgg ccc tgc act ctc ctg ttt ttt ctt ctc ttc atc cct 48
 Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
 1 5 10 15

gtc ttc tgc aaa gca atg cac gtg gcc cag cct gct gtg gta ctg gcc 96
 Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
 20 25 30

agc agc cga gcc atc gcc agc ttt gtg tgt gag tat gca tct cca gcc 144
 Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
 35 40 45

aaa gcc act gag gtc cgg gtg aca gtg ctt cgg cag gct gac agc cag 192
 Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
 50 55 60

gtg act gaa gtc tgt gcg gca acc tac atg atg ggg aat gag ttg acc 240
 Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
 65 70 75 80

ttc cta gat gat tcc atc tgc acg gcc acc tcc agt gga aat caa gtg 288
 Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
 85 90 95

aac ctc act atc caa gga ctg agg gcc atg gac acg gga ctc tac atc 336
 Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
 100 105 110

Sequence Listing

tgc aag gtg gag ctc atg tac cca ccg cca tac tac ctg ggc ata ggc Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly 115 120 125	384
aac gga acc cag att tat gta att gat cca gaa ccg tgc cca gat tcg Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser 130 135 140	432
gat aac atg cac gtg gcc cag cct gct gtg gta ctg gcc agc agc cga Asp Asn Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg 145 150 155 160	480
ggc atc gcc agc ttt gtg tgt gag tat gca tct cca ggc aaa gcc act Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr 165 170 175	528
gag gtc cgg gtg aca gtg ctt cgg cag gct gac agc cag gtg act gaa Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu 180 185 190	576
gtc tgt gcg gca acc tac atg atg ggg aat gag ttg acc ttc cta gat Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp 195 200 205	624
gat tcc atc tgc acg gcc acc tcc agt gga aat caa gtg aac ctc act Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr 210 215 220	672
atc caa gga ctg agg gcc atg gac acg gga ctc tac atc tgc aag gtg Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val 225 230 235 240	720
gag ctc atg tac cca ccg cca tac tac ctg ggc ata ggc aac gga acc Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr 245 250 255	768
cag att tat gta att gat cca gaa ccg tgc cca gat tct gca gag ccc Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Ala Glu Pro 260 265 270	816

Sequence Listing

aaa tct tgt gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu 275 280 285	864
ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp 290 295 300	912
acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg gtg gtg gac Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp 305 310 315 320	960
gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly 325 330 335	1008
gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag cag tac aac Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn 340 345 350	1056
agc acg tac cgg gtg gtc agc gtc ctc acc gtc tgt cac cag gac tgg Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His Gln Asp Trp 355 360 365	1104
ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro 370 375 380	1152
gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu 385 390 395 400	1200
cca cag gtg tac acc ctg ccc cca tcc cgg gat gag ctg acc aag aac Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn 405 410 415	1248
cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile 420 425 430	1296

Sequence Listing

gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag acc 1344
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
435 440 445

acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tac agc aag 1392
Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
450 455 460

ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc 1440
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
465 470 475 480

tcc gtg atg cat gag gct ctg cac aac cac tac acg cag aag agc ctc 1488
Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
485 490 495

tcc ctg tct ccg ggt aaa tga 1509
Ser Leu Ser Pro Gly Lys
500

<210> 20

<211> 502

<212> PRT

<213> Homo sapiens

<400> 20

Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
1 5 10 15

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
20 25 30

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
35 40 45

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
50 55 60

Sequence Listing

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
65 70 75 80

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
 85 90 95

Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
 100 105 110

Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
 115 120 125

Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser
 130 135 140

Asp Asn Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg
145 150 155 160

Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr
 165 170 175

Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu
 180 185 190

Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp
 195 200 205

Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr
 210 215 220

Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val
225 230 235 240

Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr
 245 250 255

Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Ala Glu Pro
 260 265 270

Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu

Sequence Listing

275	280	285
Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp		
290	295	300
Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp		
305	310	315
		320
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly		
325	330	335
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn		
340	345	350
Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His Gln Asp Trp		
355	360	365
Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro		
370	375	380
Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu		
385	390	395
		400
Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn		
405	410	415
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile		
420	425	430
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr		
435	440	445
Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys		
450	455	460
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys		
465	470	475
		480
Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu		
485	490	495

Sequence Listing

Ser Leu Ser Pro Gly Lys

500

<210> 21

<211> 1854

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1851)

<223> mgCD2-CD2-IgG

<220>

<221> C_region

<222> (1153)..(1854)

<223> Hinge, CH2, CH3 region

<220>

<221> misc_signal

<222> (265)..(273)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (421)..(429)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (448)..(456)

<223> N-linked glycosylation site

Sequence Listing

<220>
<221> misc_signal
<222> (598)..(606)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (616)..(624)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (805)..(813)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (961)..(969)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (988)..(996)
<223> N-linked glycosylation site

<220>
<221> primer_bind
<222> (1)..(27)
<223> PCR primer SEQ ID : 40 binding site

<220>
<221> primer_bind
<222> (588)..(630)
<223> PCR primer SEQ ID : 50(antisense) binding site

Sequence Listing

<220>
 <221> primer_bind
 <222> (588)..(630)
 <223> PCR primer SEQ ID : 49 binding site

<220>
 <221> primer_bind
 <222> (1128)..(1158)
 <223> PCR primer SEQ ID : 41(antisense) binding site

<220>
 <221> primer_bind
 <222> (1151)..(1173)
 <223> PCR primer SEQ ID : 42 binding site

<220>
 <221> primer_bind
 <222> (1832)..(1854)
 <223> PCR primer SEQ ID : 28(antisense) binding site

<220>
 <221> sig_peptide
 <222> (1)..(72)
 <223> signal peptide

<400> 21
 atg agc ttt cca tgt aaa ttt gta gcc agc ttc ctt ctg att ttc aat 48
 Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn
 1 5 10 15
 gtt tct tcc aaa ggt gca gtc tcc aaa gag att acg aat gcc ttg gaa 96
 Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu
 20 25 30

Sequence Listing

acc tgg ggt gcc ttg ggt cag gac atc aac ttg gac att cct agt ttt	144
Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe	
35 40 45	
caa atg agt gat gat att gac gat ata aaa tgg gaa aaa act tca gac	192
Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp	
50 55 60	
aag aaa aag att gca caa ttc aga aaa gag aaa gag act ttc aag gaa	240
Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu	
65 70 75 80	
aaa gat aca tat aag cta ttt aaa aat gga act ctg aaa att aag cat	288
Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His	
85 90 95	
ctg aag acc gat gat cag gat atc tac aag gta tca ata tat gat aca	336
Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr	
100 105 110	
aaa gga aaa aat gtg ttg gaa aaa ata ttt gat ttg aag att caa gag	384
Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu	
115 120 125	
agg gtc tca aaa cca aag atc tcc tgg act tgt atc aac aca acc ctg	432
Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu	
130 135 140	
acc tgt gag gta atg aat gga act gac ccc gaa tta aac ctg tat caa	480
Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln	
145 150 155 160	
gat ggg aaa cat cta aaa ctt tot cag agg gtc atc aca cac aag tgg	528
Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp	
165 170 175	
acc acc agc ctg agt gca aaa ttc aag tgc aca gca ggg aac aaa gtc	576
Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val	
180 185 190	

Sequence Listing

agc aag gaa tcc agt gtc gag aat gtc agc tgt cct aaa aat att acg	624
Ser Lys Glu Ser Ser Val Glu Asn Val Ser Cys Pro Lys Asn Ile Thr	
195 200 205	
aat gcc ttg gaa acc tgg ggt gcc ttg ggt cag gac atc aac ttg gac	672
Asn Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp	
210 215 220	
att cct agt ttt caa atg agt gat gat att gac gat ata aaa tgg gaa	720
Ile Pro Ser Phe Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu	
225 230 235 240	
aaa act tca gac aag aaa aag att gca caa ttc aga aaa gag aaa gag	768
Lys Thr Ser Asp Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu	
245 250 255	
act ttc aag gaa aaa gat aca tat aag cta ttt aaa aat gga act ctg	816
Thr Phe Lys Glu Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu	
260 265 270	
aaa att aag cat ctg aag acc gat gat cag gat atc tac aag gta tca	864
Lys Ile Lys His Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser	
275 280 285	
ata tat gat aca aaa gga aaa aat gtg ttg gaa aaa ata ttt gat ttg	912
Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu	
290 295 300	
aag att caa gag agg gtc tca aaa cca aag atc tcc tgg act tgt atc	960
Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile	
305 310 315 320	
aac aca acc ctg acc tgt gag gta atg aat gga act gac ccc gaa tta	1008
Asn Thr Thr Leu Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu	
325 330 335	
aac ctg tat caa gat ggg aaa cat cta aaa ctt tct cag agg gtc atc	1056
Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile	
340 345 350	

Sequence Listing

aca cac aag tgg acc acc agc ctg agt gca aaa ttc aag tgc aca gca Thr His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala 355 360 365	1104
ggg aac aaa gtc agc aag gaa tcc agt gtc gag cct gtc agc tgt cct Gly Asn Lys Val Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro 370 375 380	1152
gca gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg tgc cca Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro 385 390 395 400	1200
gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys 405 410 415	1248
ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val 420 425 430	1296
gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr 435 440 445	1344
gtg gac ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu 450 455 460	1392
cag tac aac agc acg tac cgg gtg gtc agc gtc ctc acc gtc tgt cac Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His 465 470 475 480	1440
cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys 485 490 495	1488
gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln 500 505 510	1536

Sequence Listing

ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat gag ctg 1584
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
515 520 525

acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc 1632
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
530 535 540

agc gac atc gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac 1680
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
545 550 555 560

tac aag acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc 1728
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
565 570 575

tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc 1776
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
580 585 590

ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg cag 1824
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
595 600 605

aag agc ctc tcc ctg tct ccg ggt aaa tga 1854
Lys Ser Leu Ser Leu Ser Pro Gly Lys
610 615

<210> 22

<211> 617

<212> PRT

<213> Homo sapiens

<400> 22

Met Ser Phe Pro Cys Lys Phe Val Ala Ser Phe Leu Leu Ile Phe Asn
1 5 10 15

Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn Ala Leu Glu

Sequence Listing

20	25	30
Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser Phe		
35	40	45
Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp		
50	55	60
Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu		
65	70	75
Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His		
85	90	95
Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr		
100	105	110
Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu Lys Ile Gln Glu		
115	120	125
Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu		
130	135	140
Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu Asn Leu Tyr Gln		
145	150	155
Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr His Lys Trp		
165	170	175
Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys Val		
180	185	190
Ser Lys Glu Ser Ser Val Glu Asn Val Ser Cys Pro Lys Asn Ile Thr		
195	200	205
Asn Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp		
210	215	220
Ile Pro Ser Phe Gln Met Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu		
225	230	235
		240

Sequence Listing

Lys Thr Ser Asp Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu
245 250 255

Thr Phe Lys Glu Lys Asp Thr Tyr Lys Leu Phe Lys Asn Gly Thr Leu
260 265 270

Lys Ile Lys His Leu Lys Thr Asp Asp Gln Asp Ile Tyr Lys Val Ser
275 280 285

Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu Lys Ile Phe Asp Leu
290 295 300

Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser Trp Thr Cys Ile
305 310 315 320

Asn Thr Thr Leu Thr Cys Glu Val Met Asn Gly Thr Asp Pro Glu Leu
325 330 335

Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile
340 345 350

Thr His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala
355 360 365

Gly Asn Lys Val Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro
370 375 380

Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
385 390 395 400

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
405 410 415

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
420 425 430

Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
435 440 445

Sequence Listing

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
450 455 460

Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His
465 470 475 480

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
485 490 495

Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
500 505 510

Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
515 520 525

Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
530 535 540

Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
545 550 555 560

Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
565 570 575

Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
580 585 590

Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
595 600 605

Lys Ser Leu Ser Leu Ser Pro Gly Lys
610 615

<210> 23
<211> 1509
<212> DNA
<213> Homo sapiens

<220>

Sequence Listing

<221> CDS
<222> (1)..(1506)
<223> mgCTLA4-CTLA4-IgG

<220>
<221> C_region
<222> (808)..(1509)
<223> Hinge, CH2, CH3 region

<220>
<221> misc_signal
<222> (289)..(297)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (385)..(393)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (403)..(411)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (424)..(432)
<223> N-linked glycosylation site

<220>
<221> misc_signal
<222> (439)..(447)
<223> N-linked glycosylation site

Sequence Listing

<220>

<221> misc_signal

<222> (664)..(672)

<223> N-linked glycosylation site

<220>

<221> misc_signal

<222> (760)..(768)

<223> N-linked glycosylation site

<220>

<221> primer_bind

<222> (1)..(15)

<223> PCR primer SEQ ID : 43 binding site

<220>

<221> primer_bind

<222> (394)..(456)

<223> PCR primer SEQ ID : 52(antisense) binding site

<220>

<221> primer_bind

<222> (397)..(460)

<223> PCR primer SEQ ID : 51 binding site

<220>

<221> primer_bind

<222> (784)..(813)

<223> PCR primer SEQ ID : 44(antisense) binding site

<220>

<221> primer_bind

<222> (805)..(826)

Sequence Listing

<223> PCR primer SEQ ID : 42 binding site

<220>

<221> primer_bind

<222> (1486)..(1509)

<223> PCR primer SEQ ID : 28(antisense) binding site

<220>

<221> sig_peptide

<222> (1)..(63)

<223> signal peptide

<400> 23

atg agg acc tgg ccc tgc act ctc ctg ttt ttt ctt ctc ttc atc cct 48

Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro

1 5 10 15

gtc ttc tgc aaa gca atg cac gtg gcc cag cct gct gtg gta ctg gcc 96

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala

20 25 30

agc agc cga ggc atc gcc agc ttt gtg tgt gag tat gca tct cca ggc 144

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly

35 40 45

aaa gcc act gag gtc cgg gtg aca gtg ctt cgg cag gct gac agc cag 192

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln

50 55 60

gtg act gaa gtc tgt gcg gca acc tac atg atg ggg aat gag ttg acc 240

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr

65 70 75 80

ttc cta gat gat tcc atc tgc acg ggc acc tcc agt gga aat caa gtg 288

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val

85 90 95

Sequence Listing

aac ctc act atc caa gga ctg agg gcc atg gac acg gga ctc tac atc Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile 100 105 110	336
tgc aag gtg gag ctc atg tac cca ccg cca tac tac ctg ggc ata ggc Cys Lys Val Glu Leu Met Tyr Pro Pro Tyr Tyr Leu Gly Ile Gly 115 120 125	384
aac gga acc cag att tat gta aat gat aca gaa ccg tgc aat gat tcg Asn Gly Thr Gln Ile Tyr Val Asn Asp Thr Glu Pro Cys Asn Asp Ser 130 135 140	432
gat aac aat cac acg gcc cag cct gct gtg gta ctg gcc agc agc cga Asp Asn Asn His Thr Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg 145 150 155 160	480
ggc atc gcc agc ttt gtg tgt gag tat gca tct cca ggc aaa gcc act Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr 165 170 175	528
gag gtc cgg gtg aca gtg ctt cgg cag gct gac agc cag gtg act gaa Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu 180 185 190	576
gtc tgt gcg gca acc tac atg atg ggg aat gag ttg acc ttc cta gat Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp 195 200 205	624
gat tcc atc tgc acg ggc acc tcc agt gga aat caa gtg aac ctc act Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr 210 215 220	672
atc caa gga ctg agg gcc atg gac acg gga ctc tac atc tgc aag gtg Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val 225 230 235 240	720
gag ctc atg tac cca ccg cca tac tac ctg ggc ata ggc aac gga acc Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr 245 250 255	768

Sequence Listing

cag att tat gta att gat cca gaa ccg tgc cca gat tct gca gag ccc Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Ala Glu Pro 260 265 270	816
aaa tct tgt gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu 275 280 285	864
ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp 290 295 300	912
acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg gtg gtg gac Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp 305 310 315 320	960
gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly 325 330 335	1008
gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag cag tac aac Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn 340 345 350	1056
agc acg tac cgg gtg gtc agc gtc ctc acc gtc tgt cac cag gac tgg Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His Gln Asp Trp 355 360 365	1104
ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro 370 375 380	1152
gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu 385 390 395 400	1200
cca cag gtg tac acc ctg ccc cca tcc cgg gat gag ctg acc aag aac Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn 405 410 415	1248

Sequence Listing

cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc 1296
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
420 425 430

gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag acc 1344
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
435 440 445

acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tac agc aag 1392
Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
450 455 460

ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc 1440
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
465 470 475 480

tcc gtg atg cat gag gct ctg cac aac cac tac acg cag aag agc ctc 1488
Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
485 490 495

tcc ctg tct ccg ggt aaa tga 1509
Ser Leu Ser Pro Gly Lys
500

<210> 24

<211> 502

<212> PRT

<213> Homo sapiens

<400> 24

Met Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
1 5 10 15

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
20 25 30

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
35 40 45

Sequence Listing

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
50 55 60

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
65 70 75 80

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
85 90 95

Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
100 105 110

Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
115 120 125

Asn Gly Thr Gln Ile Tyr Val Asn Asp Thr Glu Pro Cys Asn Asp Ser
130 135 140

Asp Asn Asn His Thr Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg
145 150 155 160

Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr
165 170 175

Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu
180 185 190

Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp
195 200 205

Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr
210 215 220

Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val
225 230 235 240

Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr
245 250 255

Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Ala Glu Pro

Sequence Listing

260	265	270
Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu		
275	280	285
Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp		
290	295	300
Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp		
305	310	315 320
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly		
325	330	335
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn		
340	345	350
Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Cys His Gln Asp Trp		
355	360	365
Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro		
370	375	380
Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu		
385	390	395 400
Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn		
405	410	415
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile		
420	425	430
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr		
435	440	445
Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys		
450	455	460
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys		
465	470	475 480

Sequence Listing

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
485 490 495

Ser Leu Ser Pro Gly Lys
500

<210> 25
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide TNFR1-EDF-EcoRI

<400> 25
ccggaattcc ggtctggcat ggcctctcc acc 33

<210> 26
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide TNFR1-EDR-IgGh

<400> 26
cacaagattt ggcctctgct gtggtgctg agtcctc 37

<210> 27
<211> 37
<212> DNA
<213> Artificial Sequence

<220>

Sequence Listing

<223> PCR primer, oligonucleotide IgG1-T1F

<400> 27

gaggactcag gcaccacagc agagcccaaa tttgtg

37

<210> 28

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide IgG1-R-XbaI

<400> 28

gctctagagc tcatttaccc ggagacaggg agag

34

<210> 29

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide TNFR2-EDF-EcoRI

<400> 29

cgggaattcc gggcaccocat ggcgccgto gcc

33

<210> 30

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide TNFR2-EDR-IgGh

Sequence Listing

<400> 30

cacaagattt gggctctgcg tcgccagtgc tcccttc

37

<210> 31

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide IgG-T2F

<400> 31

gaagggagca ctggcgacgc agagcccaaa tcttgtg

37

<210> 32

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide TNFR1-CF-BamHI

<400> 32

cgcggatccg ggaacatttc actggtecct cacctag

37

<210> 33

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide TNFR1-NR-BamHI

Sequence Listing

<400> 33
cgcggatccg tcctcagtcg ccttaacatt ctcaatctg 39

<210> 34
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide TNFR2-CF-BamHI

<400> 34
cgcggatcca acgcaactac accctacgac cggag 36

<210> 35
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide TNFR2-NR-BamHI

<400> 35
cgcggatccg ctcccttcag ctggggggct g 31

<210> 36
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide mgTNFR1-TNFR1-IgG-F

Sequence Listing

<400> 36
aaaagcaacg agaccaacaa gacctgccta cacaacgggt ccaggagagaa gaacgatagt 60

gtg 63

<210> 37
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide mgTNFR1-TNFR1-IgG-R

<400> 37
ciccctggac ccgttggtga ggcagggtctt gttggtctcg ttgcttttct tacagttact 60

ac 62

<210> 38
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer, oligonucleotide mgTNFR2-TNFR2-IgG-F

<400> 38
atggatgcaa actgcacgtc cccggagccc aacagcacat gccgg 45

<210> 39
<211> 42
<212> DNA
<213> Artificial Sequence

<220>

Sequence Listing

<223> PCR primer, oligonucleotide mgTNFR2-TNFR2-IgG-R

<400> 39

gcatgtgctg ttgggtccg gggacgtgca gtttgcaccc at

42

<210> 40

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CD2F-EcoRI

<400> 40

ccggaattca tgagctttcc atgtaaattt gtagcc

36

<210> 41

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CD2R-PstI

<400> 41

ctctgcagga cagctgacag gctcgacact

30

<210> 42

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide IgG-F-PstI

Sequence Listing

<400> 42

atctgcagag cccaaatctt gtgac

25

<210> 43

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CTLA4F-EcoRI

<400> 43

cgggaattca tgaggacctg gcc

24

<210> 44

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CTLA4R-PstI

<400> 44

ctctgcagaa tctgggcacg gttcaggatc

30

<210> 45

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CD2-NT-F

Sequence Listing

<400> 45

taaagagatt acgaatgcc

19

<210> 46

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CD2-CT-R

<400> 46

tgacaggacag ctgacagg

18

<210> 47

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CTLA4-NT-F

<400> 47

ggataatcat gcacgtggcc cag

23

<210> 48

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide CTLA4-CT-R

Sequence Listing

<400> 48

tgcagaatct gggcacgg

18

<210> 49

<211> 43

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide mgCD2-CD2-IgG-F

<400> 49

cagtgtcgag aatgtcagct gtcctaaaaa tattacgaat gcc

43

<210> 50

<211> 43

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide mgCD2-CD2-IgG-R

<400> 50

ggcattcgta atatttttag gacagctgac attctcgaca ctg

43

<210> 51

<211> 64

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide mgCTLA4-CTLA4-IgG-F

<400> 51

Sequence Listing

atttatgtaa acgatacaga accgtgcaat gattcggata acaaccacac agcccagcct 60

gctg 64

<210> 52

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer, oligonucleotide mgCTLA4-CTLA4-IgG-R

<400> 52

aggctgggct gtgtggttgt taccgaatc attgcacggt tctgtatcgt ttacataaat 60

ctg 63

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR02/01427

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C07K 16/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C07K 16/46, C07K 19/00, C12N 15

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean Patents and applications for inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Medline, Biosis

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,073,627 A (Immunex Corporation) 17 DECEMBER 1991 see the whole document	1
X, P ----- Y, P	EP1148065 A1 (ROSE-JOHN, STEFAN) 24 OCTOBER 2001 see column3, lines 20-40, claims	1 --- 2-5, 7-10, 12, 14, 15
Y	EP0464533 A1 (HOECHST AKTIENGESELLSCHAFT) 8 JANUARY 1992 see claims	2-5, 7-10, 12, 14, 15
Y	US 5861151 A (BRISTOL-MYERS SQUIBB CO.) 19 JANUARY 1999 see column7, lines 40-45, Fig.1	2-5, 7-10, 12, 14, 15
A	US 5349053 A (PROTEIN DESIGN LABS, INC) 20 SEPTEMBER 1994 see the whole document	1-35
A	US 5428130 A (GENENTECH, INC) 27 JUNE 1995 see the whole document	1-35
A	US 6165476 A (BETH ISRAEL DEACONESS MEDICAL) 26 DECEMBER 2000 see the whole document	1-35

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 DECEMBER 2002 (11.12.2002)

Date of mailing of the international search report

12 DECEMBER 2002 (12.12.2002)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

HAN, Hyun Sook

Telephone No. 82-42-481-5596



Form PCT/ISA 210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR02/01427

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5073627 A	17.12.91	AU 6424090 A1 EP 0489116 B1 WO9102754 A1	03.04.91 06.04.94 07.03.91
EP1148065 A1	24.10.01	NONE	
EP 0464533 A1	08.01.92	JP 5247094 A2 KR 0249572 B1 US 20010053539 A1	24.09.93 15.03.00 20.12.01
US 5861151 A1	19.01.99	AU 03327293 A1 EP 0619843 A1 WO 9313210 A1	28.07.93 19.10.94 19.01.99
US 5349053 A1	20.09.94	NONE	
US 5428130 A1	27.06.95	EP 1029870 A2 JP 5503009 T2 WO 9108298 A2	23.08.00 27.03.93 13.06.91
US 6165476 A1	26.12.00	AU 8392198 A1 JP 2001510682 T2 WO 9902711 A3	08.02.99 07.08.01 02.09.99